

2015 HAWAII STD SCREENING RECOMMENDATIONS

The following recommendations are based on guidelines for 2015 STD screening from the Centers for Disease Control and Prevention, United States Preventive Services Task Force, Infectious Disease Society of America, and the Hawaii STD/AIDS Prevention Program. In populations for whom no recommendations exist, screening should be based on risk factors, local epidemiology and prevalence of specific STDs in the particular clinical setting. **All women** diagnosed with chlamydia, gonorrhea, or trichomonas should be retested for repeat infection at 3 months after treatment. Men diagnosed with chlamydia (CT) or gonorrhea (GC) should also be retested at 3 months. Retesting can also be performed opportunistically anytime the patient returns for care in the 1-12 months after treatment. Other factors to consider prior to screening are summarized in the footnotes below. Call the Hawaii STD/AIDS Prevention Program at (808) 733-9281 for queries. **DISCLAIMER:** This does not reflect an exhaustive list. Refer to <http://www.cdc.gov/std/tg2015/screening-recommendations.htm> for more complete information.

	Population	STD Screening Recommendations	Frequency	Comments
Women	Women < 25 years of age ^{1,4}	CT and GC (vaginal, cervical, or urine)..... Other STDs according to risk. HIV	Annually At least once, then repeat according to level of risk	CT/GC: consider screening more frequently for those at increased risk.
	Women 25 years of age and older ^{1,4}	No routine screening for STDs. Screen according to risk. HIV	All women up to 64 years of age at least once, then repeat according to level of risk	Targeted CT/GC screening recommended for women with risk factors. See footnote 4.
	Pregnant women ^{1,3, 5}	CT and GC (vaginal, cervical, or urine)..... Syphilis..... HIV	First trimester First trimester First trimester	Repeat screening for CT, GC, HIV, HBsAg in third trimester if at increased risk. Repeat syphilis screening recommended in third trimester if high risk or living in area of high morbidity, per local guidelines.
	HIV-positive women ^{1,6-8}	CT and GC (vaginal, cervical, or urine)..... CT and GC (rectal) GC (pharyngeal) Syphilis..... Trichomoniasis	Annually Annually (if exposed) Annually (if exposed) Annually Annually Repeat screening every 3-6 months, if indicated by risk	
		Hepatitis B Surface Antigen (HBsAg)..... Hepatitis C.....	First visit First visit	
Men	Heterosexual men ³	No routine screening for STDs. Screen according to risk. HIV	All men 13-64 years of age at least once, then annually if high-risk	Targeted screening for CT in high risk settings (e.g. corrections) or if risk factors (e.g. CT in past 24 months)
	Men who have sex with men (MSM) ^{1,6}	CT and GC (urine) CT and GC (rectal) GC (pharyngeal) Syphilis..... HIV	Annually Annually (if exposed) Annually (if exposed) Annually Annually Repeat screening every 3-6 months, as indicated by risk	
		Hepatitis B Surface Antigen (HBsAg).....	At least once	
	HIV-positive men ⁶⁻⁸	CT and GC (urine) CT and GC (rectal) GC, (pharyngeal) Syphilis.....	Annually Annually (if exposed) Annually (if exposed) Annually Repeat screening every 3-6 months, as indicated by risk	
		Hepatitis B Surface Antigen (HBsAg)..... Hepatitis C.....	First visit Annually	

¹CDC. STD Treatment Guidelines. June 5, 2015 / 64(RR3);1-137.

²Human papillomavirus (HPV) testing is recommended as part of cervical cancer screening and management of cervical intraepithelial neoplasia. It is not recommended as part of routine STD screening or prior to initiating HPV vaccination. See the American Society for Colposcopy and Cervical Pathology (www.asccp.org) for further guidance.

³Screening for asymptomatic HSV-2 infection should be offered to select patients including those in partnerships or considering partnerships with HSV-2-infected individuals. Counseling should be provided to patients tested for HSV-2.

⁴Risk factors for CT or GC: prior CT or GC infection, particularly in past 24 months; more than one sex partner in the past year; suspicion that a recent partner may have had concurrent partners; new sex partner in past 3 months; exchanging sex for drugs or money in the past year; and local factors such as community prevalence of infection.

⁵In pregnant women with a history of injection drug use or a history of blood transfusion or organ transplantation before 1992, screening for hepatitis C should be conducted

⁶Routine hepatitis B vaccination is recommended for all HIV-infected patients and all MSM. Routine hepatitis A vaccination is recommended for all MSM. Pre-vaccination serologic testing may be considered, however if testing is not feasible in the current setting, routine vaccination should continue.

⁷Primary Care Guidelines for the Management of Persons Infected with Human Immunodeficiency Virus: 2013 Update by the HIV Medicine Association of the Infectious Disease Society of America. *Clinical Infectious Diseases* 2013; doi: 10.1093/cid/cit665.

⁸Data are insufficient to recommend routine anal cancer screening with anal cytology among HIV-positive men and women. Some clinical centers perform anal cytology screening in populations at high-risk for anal cancer. Programmatic considerations such as availability of providers to perform diagnostic anoscopy in the case of abnormal results should be considered prior to initiating anal cancer screening.