TABLE 63 INFANT DEATHS BY GEOGRAPHICAL AREA 2000

	INFANT DEATHS				PERINATAL MORTALITY	
	PLACE OF OCCURRENCE PLACE OF RESIDENCE		RESIDENCE	PLACE OF RESIDENCE		
COUNTY	Number	Rate <u>a</u> /	Number	Rate <u>a</u> /	Number	Rate <u>b</u> /
Hawai'i	*	4.2	*	10.3	*	12.8
Honolulu	*	9.2	*	7.5	*	8.7
Kaua'i	*	2.7	*	2.7	*	9.3
Maui	*	3.5	*	8.0	*	8.0
Sub-Total Residents	-	-	*	7.7	*	9.1
Non-Residents	-	-	*	-	*	-
Unknown	-	-	*	-	*	-
TOTAL	138	7.8	138	-	173	9.8

^{*}Cell sizes suppressed.

TABLE 64
INFANT DEATHS BY ETHNICITY OF INFANT, AGE AND BIRTH WEIGHT RESIDENTS, 2000

	AGE OF INFANT (DAYS)						BIRTH WEIGHT
ETHNICITY	Under 1 Day	1 - 6	7 - 27	Over 27	Total		Under 2500
OF INFANT	Rate <u>a</u> /	Rate <u>a</u> /	Rate <u>a</u> /	Rate <u>a</u> /	No.	Rate <u>a</u> /	grams <u>b</u> /
Caucasian	3.0	2.0	1.3	1.3	23	7.7	*
Hawaiian	3.7	1.0	8.0	2.6	50	8.0	31
Filipino	2.7	1.7	2.0	1.3	23	7.7	16
Japanese	2.1	1.6	0.5	1.6	11	5.8	*
All Others	2.4	2.1	2.1	1.5	27	8.0	17
TOTAL	3.0	1.5	1.3	1.8	134	7.7	87

^{*}Cell size suppressed because one or more cells within the column is less than 5.

Notes:

a/ Number of Resident Deaths per 1,000 live births.

b/ Resident infant deaths under 1 week of age plus the number of fetal deaths of 20 weeks or more gestation divided by the sum of total resident live births plus resident fetal deaths of 20 weeks or more gestation, per 1,000.

a/ Rates are number of infant deaths per 1,000 live births.

b/ Birth weights under 2,500 grams are considered low birth weight.

TABLE 65 INFANT DEATHS BY CAUSE, SEX, BIRTH WEIGHT AND AGE RESIDENTS, 2000

		NEONATAL <u>a</u> /		
CAUSE OF DEATH <u>b</u> /	Number of Infant Deaths	Under 28 Days	Rate	
Diseases of the Circulatory System	7	5	0.3	
Maternal Complication of Pregnancy	12	12	0.7	
Complications of Placenta, Cord & Membrane	9	9	0.5	
Short Gestation	12	12	0.7	
Hypoxia/Asphyxia	6	5	0.3	
Other Respiratory Conditions	14	11	0.6	
Perinatal Infections	6	6	0.3	
Other Perinatal Conditions	12	11	0.6	
Congenital Anomaly	20	15	0.9	
All Other Causes	36	16	0.9	
TOTAL	134	102	5.8	

^{*}Cell size suppressed.

Note:

a/ Neonatal deaths are deaths of infants under 28 days of age. Rate is per 1,000 live births.

b/ The method of coding causes of death changed in 1999 to the International Classification of Diseases, Version 10.