

State of Hawaii SARS-CoV-2 Sequencing and Variant Report

Hawaii Department of Health

2022-06-22

Contents

Introduction	3
Acknowledgements	4
County distribution of genomes sequenced by State Laboratories Division since Jan 1st, 2021	4
Summary and key notes	5
Variants of Concern (VOC) and Variants Being Monitored (VBM)	6
Variants of Concern that have been detected in the State of Hawaii	6
Omicron variant (B.1.1.529, BA.* lineages, and XE)	6
Variants Being Monitored that have been detected in the State of Hawaii	6
Alpha variant (B.1.1.7 and Q.* lineages)	6
Beta variant (B.1.351 and B.1.351.* lineages)	6
Gamma variant (P.1 and P.1.* lineages)	7
Delta variant (B.1.617.2 and AY.* lineages)	7
Epsilon variant (B.1.429 and B.1.427)	7
Zeta variant (P.2)	7
Iota variant (B.1.526)	7
Mu variant (B.1.621 and B.1.621.1)	7
State of Hawaii	8
Total variants identified	8
Estimated proportions of variants circulating in the State of Hawaii	9
Variants of Concern in the State of Hawaii	10
Variants Being Monitored in the State of Hawaii	11
Honolulu County	12
Total variants identified	12
Estimated proportions of variants circulating in Honolulu County	13
Variants of Concern in Honolulu County	14
Variants Being Monitored in Honolulu County	15
Maui County	16
Total variants identified	16
Estimated proportions of variants circulating in Maui County	17
Variants of Concern in Maui County	18
Variants Being Monitored in Maui County	19
Hawaii County	20
Total variants identified	20
Estimated proportions of variants circulating in Hawaii County	21
Variants of Concern in Hawaii County	22
Variants Being Monitored in Hawaii County	23
Kauai County	24
Total variants identified	24
Estimated proportions of variants circulating in Kauai County	25
Variants of Concern in Kauai County	26
Variants Being Monitored in Kauai County	27

Introduction

Whole genome sequencing (WGS) involves a set of laboratory methods used to determine the full genome sequence of an organism or virus, which in the case of Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2), the virus that causes Coronavirus Disease 2019 (COVID-19), is approximately 30,000 letters, each letter a nucleotide code denoted as A, C, G, or T.

The genome sequence of a virus can reveal mutations that make it unique. Mutations are changes in a genome sequence (usually one-letter changes) that occur naturally over time.

Collecting the genome sequences of virus specimens can reveal information about the relatedness of viruses and the similarities shared among groups of viruses. Groups of same-species viruses that share a set of genome mutations are referred to as a lineage.

Scientists compare viral genomes to better understand virus transmission, how viruses can spread from person to person. Sequencing also allows public health officials to monitor viruses involved in outbreaks, characterize outbreaks, detect clusters of cases, and monitor new lineages. Novel mutations can emerge with new lineages and scientists refer to these new lineages as emerging variants.

Some of these variants are classified by the Centers for Disease Control and Prevention (CDC) as a Variant of Concern (VOC) and others as Variants Being Monitored (VBM), because of their attributes, which, for example, can be increased transmissibility, decreased neutralization by antibodies generated during previous infection or vaccination, and/or increased severity of disease. The CDC has extensive information about SARS-CoV-2 variant classification (<https://www.cdc.gov/coronavirus/2019-ncov/cases-updates/variant-surveillance/variant-info.html>), which is updated as new evidence becomes available.

Sequencing can only be performed on specimens that contain SARS-CoV-2 RNA, which means only specimens used for molecular tests (such as PCR) can be included. Therefore, this report is limited to confirmed PCR-positives only. The genomes that are sequenced and compared are those of the virus, not humans.

Sequencing can be performed on stored specimens at any time. Therefore, the dataset used for this report is dynamic and batches of stored specimens that are newly sequenced will be added to the dataset as sequencing occurs. Because of this, trends based on historical data can change over time.

In February 2021, State Laboratories Division, Hawaii Department of Health increased sequencing efforts done on PCR-positive specimens to improve the State's ability to detect new variants. According to the CDC, approximately 5% of PCR-positive cases in the State of Hawaii have been sequenced since testing began (<https://covid.cdc.gov/covid-data-tracker/#published-sars-cov-2-sequences>).

Acknowledgements

This report integrates genomes sequenced since Jan 1, 2021 by:

Institution	Program/partner	Count	Percent
Laboratory Preparedness and Response Branch, State Laboratories Division		10497	74.310%
Centers for Disease Control and Prevention	National SARS-CoV-2 Strain Surveillance (NS3)	412	2.917%
	Quest Diagnostics Incorporated	958	6.782%
	Laboratory Corporation of America	1036	7.334%
	Aegis Sciences Corporation	379	2.683%
	Helix/Illumina	119	0.842%
	Infinity Biologix	16	0.113%
	Mako Medical	20	0.142%
	Fulgent Genetics	8	0.057%
Tripler Army Medical Center		681	4.821%
Total		14126	100.000%

Table Notes:

- In support of the State’s SARS-CoV-2 genomic surveillance program, PCR-positive specimens are regularly provided to the Laboratory Preparedness and Response Branch (LPRB), State Laboratories Division by the various commercial and medical laboratories that operate throughout the State of Hawaii, to include Clinical Labs of Hawaii (CLH), Diagnostic Laboratory Services, Inc. (DLS), Kaiser Permanente Hawaii, and S&G Labs Hawaii. The collected specimens are then quality controlled and sequenced by the LPRB.
- The specimens sequenced by the Centers for Disease Control and Prevention (CDC) for the National SARS-CoV-2 Strain Surveillance (NS3) program are collected, quality controlled, and shipped to the CDC by the LPRB.

County distribution of genomes sequenced by State Laboratories Division since Jan 1st, 2021

Honolulu County	Maui County	Hawaii County	Kauai County	unknown	Total
6222	1701	1686	546	342	10497

Table Notes:

- County information is unavailable or “unknown” for a number of specimens sequenced by State Laboratories Division.
- County information is not provided for specimens sequenced by the CDC and its commercial partners, and by Tripler Army Medical Center (TAMC).

Summary and key notes

- The data in this report reflect sequenced PCR-positive specimens only, and not all PCR-positive cases in the State of Hawaii are sequenced.
- Each successfully sequenced specimen produces one consensus SARS-CoV-2 genome sequence that is further analyzed to determine the variant.
- State Laboratories Division has reported 557 additional SARS-CoV-2 genome sequences since the previous report was generated (6/8/2022).
- The CDC and its commercial partners have reported 177 additional SARS-CoV-2 genome sequences from the State of Hawaii since the previous report was generated (6/8/2022).
- TAMC has not reported any additional SARS-CoV-2 genome sequences from the State of Hawaii since the previous report was generated (6/8/2022).
- SARS-CoV-2 variant nomenclature is defined by a World Health Organization (WHO) label (letters of the Greek Alphabet, i.e., Alpha, Beta, Gamma, Delta, etc.), Pango lineage (alphabetical prefix and a numerical suffix), and/or Nextstrain clade (year of emergence followed by the next available letter in the alphabet, i.e., 20A, 20B, etc.). In this report, variant counts are reported using the WHO label and Pango lineage nomenclatures only.
- Omicron represents 100% of the variants currently circulating in the State of Hawaii.
- Classifications of Omicron lineages are in flux. The parent lineage of Omicron is B.1.1.529; sub-lineages of Omicron have the BA.* designation (i.e., BA.1, BA.2, etc.). Omicron XE, which has been detected in the State of Hawaii, is a recombinant lineage containing genetic material from BA.1 and BA.2.
- BA.2.12.1 (a sub-lineage of BA.2) is currently the dominant lineage circulating in the State of Hawaii. BA.2.12.1 contains 2 additional amino acid mutations (L452Q and S704L) in the spike protein compared to BA.2. Relative growth modeling based on genomic surveillance within the U.S. estimates BA.2.12.1 to be approximately 1.3x more transmissible than BA.2 (<https://cov-spectrum.org/explore/United%20States/Surveillance/Past3M/variants?pangoLineage=BA.2&pangoLineage1=BA.2.12.1&analysisMode=CompareToBaseline&>).
- Cases of BA.4 and BA.5, which are on the rise nationally, have been detected in the State of Hawaii. Within the spike protein alone, BA.4 and BA.5 contain 4 additional amino acid mutations (H69-, V70-, L452R, and F486V) and one reversion (R493Q) compared to BA.2. Relative growth modeling based on genomic surveillance within the U.S. estimates BA.4 and BA.5 to be approximately 1.6x and 1.7x more transmissible, respectively, than BA.2 (<https://cov-spectrum.org/explore/United%20States/Surveillance/Past3M/variants?pangoLineage=BA.2&pangoLineage1=BA.4&analysisMode=CompareToBaseline&> and <https://cov-spectrum.org/explore/United%20States/Surveillance/Past3M/variants?pangoLineage=BA.2&pangoLineage1=BA.5&analysisMode=CompareToBaseline&>).
- Preliminary reports suggest that, compared to BA.2, BA.2.12.1 and BA.4/5 have an increased ability to evade therapeutic antibodies and antibodies elicited by vaccination and/or prior infection (<https://www.biorxiv.org/content/10.1101/2022.04.30.489997v1.full>, <https://www.biorxiv.org/content/10.1101/2022.05.21.492554v1.full>, <https://www.biorxiv.org/content/10.1101/2022.05.26.493517v1.full>).

Variants of Concern (VOC) and Variants Being Monitored (VBM)

It is important to note that evidence to date shows that vaccination leads to milder cases, not requiring hospitalization, for all Variants of Concern and Variants Being Monitored that are described here, even if the efficacy of antibodies is diminished against some of these variants compared to the original version of the virus.

Also, *none* of these variants are classified as a “*Variant of High Consequence (VOHC)*”, according to CDC variant categories (<https://www.cdc.gov/coronavirus/2019-ncov/variants/variant-info.html#Consequence>), which is a category that would imply a variant has the ability to evade diagnosis, significantly reduce the vaccines’ effectiveness and protection against severe disease, significantly reduce susceptibility to treatments, or lead to more severe disease and increased hospitalizations.

Variants of Concern that have been detected in the State of Hawaii

Omicron variant (B.1.1.529, BA.* lineages, and XE)

B.1.1.529 is the parent lineage of Omicron; sub-lineages of Omicron have the BA.* designation (i.e., BA.1, BA.2, BA.4, etc.). XE is a recombinant lineage of BA.1 and BA.2, with the majority of its genome, including the gene encoding for the spike protein, coming from BA.2. SARS-CoV-2 recombinants that receive a Pango classification have an X* lineage designation (i.e., XE, XG, XH, etc.). Technical information regarding the lineage designation of XE can be found at <https://github.com/cov-lineages/pango-designation/issues/454>.

B.1.1.529 was reported to the WHO on November 24, 2021 and first detected in specimens collected on November 11, 2021 in Botswana and on November 14, 2021 in South Africa. The WHO labeled B.1.1.529 “Omicron” and classified it as a VOC on November 26, 2021. The United States designated Omicron as a VOC on November 30, 2021 and reported its first case on December 1, 2021. Omicron contains more changes in the spike protein than have been observed in other variants, including at least 30 amino acid substitutions. Several of these mutations, including S477N, N501Y, and E484K, have been associated with increased infectivity and decreased neutralizing activity of monoclonal antibodies and convalescent sera. Evidence indicates that Omicron spreads more easily than the original SARS-CoV-2 virus and the Delta variant. Lineages of Omicron also correspond to Nextstrain clades 21M, 21K, 21L, 22A, 22B, and 22C. More information about Omicron can be found at <https://www.cdc.gov/coronavirus/2019-ncov/variants/omicron-variant.html>.

Variants Being Monitored that have been detected in the State of Hawaii

Alpha variant (B.1.1.7 and Q.* lineages)

B.1.1.7 was first identified in the United Kingdom and the WHO labeled it “Alpha” on May 31, 2021. This variant contains the N501Y mutation and a short deletion in the spike protein. This variant is concerning because it has been shown to be significantly more transmissible (~50%) than the original SARS-CoV-2 lineages and reports from the United Kingdom suggest that B.1.1.7 cases are more likely to require hospitalization. B.1.1.7 does not appear to evade vaccine-induced neutralizing antibody responses. The Alpha variant also corresponds to Nextstrain clade 20I.

Beta variant (B.1.351 and B.1.351.* lineages)

B.1.351 was first identified in South Africa and the WHO labeled it “Beta” on May 31, 2021. This variant is highly infectious and can quickly spread from person to person. Preliminary studies suggest that antibodies from previous infection or from vaccination may be less effective at preventing infection against this variant due to presence of the E484K mutation in the spike protein. The Beta variant also corresponds to Nextstrain clade 20H.

Gamma variant (P.1 and P.1.* lineages)

P.1 was first identified in Brazil and the WHO labeled it “Gamma” on May 31, 2021. This variant also contains the N501Y mutation, like B.1.1.7, but not the deletion in the spike protein. Preliminary studies suggest that antibodies from previous infection or from vaccination may be less effective at preventing infection against this variant. The Gamma variant also corresponds to Nextstrain clade 20J.

Delta variant (B.1.617.2 and AY.* lineages)

B.1.617.2 is the parent lineage of Delta; sub-lineages of Delta have the AY.* designation (i.e., AY.1, AY.2, AY.3, etc.).

B.1.617.2 was first identified in India and the WHO labeled it “Delta” on May 31, 2021. This variant contains the L452R mutation in the spike protein, which has been shown to escape neutralization by monoclonal antibodies and some convalescent sera, as well as a few additional spike mutations predicted to have functional consequences (e.g. T478K). The Delta variant is highly contagious, more than 2x as contagious as previous variants. Lineages of Delta also correspond to Nextstrain clades 21A, 21I, and 21J.

Epsilon variant (B.1.429 and B.1.427)

The closely related lineages, B.1.429 and B.1.427, were first identified in California and designated initially as CA VUI1. The WHO labeled them “Epsilon” on May 31, 2021. They can quickly spread from person-to-person, with an estimated ~20% higher efficiency than the original virus. The Epsilon variant also corresponds to Nextstrain clade 21C.

Zeta variant (P.2)

P.2 was first identified in Brazil and contains a spike mutation (E484K), which is also present in B.1.351, that can potentially make it less responsive to antibodies. The Zeta variant also corresponds to Nextstrain clade 20B/S.484K.

Iota variant (B.1.526)

B.1.526 was first identified in New York and is classified by CDC as a VBM because of indications that it has increased transmissibility. The WHO labeled it “Iota” on May 31, 2021. Some of the genomes (but not all) of this variant contain the E484K mutation. The Iota variant also corresponds to Nextstrain clade 21F.

Mu variant (B.1.621 and B.1.621.1)

Lineage B.1.621 was first identified in Columbia in January 2021 and it has a couple of mutations in common with the Beta (B.1.351) and Gamma (P.1) variants, which have been associated with high transmissibility (N501Y) and a level of decreased vaccine efficiency (E484K). The MU variant also corresponds to Nextstrain clade 21H.

State of Hawaii

Total variants identified




Figure Notes:

- The graph shows the total number of variants by lineage detected in the State of Hawaii in each 2-week interval ending on the date shown (date represents when the specimen was collected from a patient).
- Omicron (B.1.1.529 + BA.* + XE) is a *Variant of Concern*.
- Delta (B.1.617.2 + AY.*) is a *Variant Being Monitored*.
- The gray bar graph (top) shows the percentage of PCR-positive cases from each 2-week time interval that were sequenced.
- The color bar graph (bottom) shows the number of viruses sequenced (one virus equates to one PCR-positive case) from each 2-week time interval.
- SARS-CoV-2 genome sequencing may not be a random sample of all cases. This graph does not estimate prevalence in the population.
- Sequencing can be performed on stored patient specimens at any time, so these numbers may change as additional specimens are sequenced.

Estimated proportions of variants circulating in the State of Hawaii




Figure Notes:

- The graph shows biweekly percentage estimates of SARS-CoV-2 variants circulating in the State of Hawaii, grouped in two-week intervals (based on the date of specimen collection).
- Not all positive SARS-CoV-2 specimens are sequenced and sequenced specimens are not a random selection of all COVID-19 cases in the State of Hawaii. This graph has been generated only counting specimens that were selected randomly for the purpose of surveillance, to avoid over-representing the specimens that were selected for investigations of clusters.
- The last 2-week interval numbers will most likely change, as a number of specimens that are currently being processed will be added.
- Sequencing of certain specimens can be delayed for technical reasons. Therefore, the dataset used for this report is dynamic and specimens that are newly sequenced will be added to the dataset as sequencing occurs. Because of this, trends based on historical data can change over time.
- Omicron (BA.1*) is an aggregation of BA.1 and its sub-lineages (i.e., BA.1.1, BA.1.15, etc.).
- Omicron (BA.2*) is an aggregation of BA.2, its sub-lineages (i.e., BA.2.3, BA.2.9, etc.), and XE; excludes BA.2.12.1.

Variants of Concern in the State of Hawaii

Variant	Lineage	Area of emergence	Cumulative cases detected	Earliest specimen collection date	Most recent specimen collection date
Omicron			7034		
	BA.1.1	South Africa/Botswana	1738	27 Nov 2021	25 May 2022
	BA.2	South Africa/Botswana	1408	18 Jan 2022	04 Jun 2022
	BA.2.3	Philippines	1021	06 Jan 2022	02 Jun 2022
	BA.2.12.1	United States/Canada	992	21 Mar 2022	04 Jun 2022
	BA.1.1.2	Japan	569	06 Dec 2021	01 Apr 2022
	BA.1.15	United States	426	30 Nov 2021	09 Apr 2022
	BA.1	South Africa/Botswana	187	07 Dec 2021	11 Apr 2022
	BA.2.9	Europe	129	25 Jan 2022	01 Jun 2022
	Other BA.*	Various	115	02 Apr 2022	May 2022
	BA.1.1.18	United States	95	10 Dec 2021	30 Mar 2022
	BA.2.18	United Kingdom	71	21 Mar 2022	03 Jun 2022
	BA.1.20	United States	44	14 Dec 2021	28 Feb 2022
	BA.2.10.1	Singapore	27	01 Feb 2022	15 May 2022
	BA.1.17.2	Europe	26	14 Dec 2021	25 Feb 2022
	BA.2.13	Europe	21	18 Apr 2022	24 May 2022
	BA.2.10	Asia	20	22 Feb 2022	24 May 2022
	BA.2.29	Japan	20	27 Mar 2022	12 May 2022
	BA.2.31	Israel	20	18 Apr 2022	29 May 2022
	BA.2.7	United States	19	28 Mar 2022	21 May 2022
	BA.1.18	Europe/North America	18	18 Dec 2021	02 Mar 2022
	BA.1.1.16	Canada/United States	17	20 Jan 2022	30 Mar 2022
	BA.2.38	India	16	21 Mar 2022	13 May 2022
	BA.5	South Africa	15	03 May 2022	03 Jun 2022
	XE	United Kingdom	11	26 Mar 2022	31 May 2022
	BA.4	South Africa	9	04 May 2022	27 May 2022

Table Notes:

- Lineage “Other BA.*” represents an aggregation of different BA.* lineages, except BA.4, in which each alone accounts for <0.2% of the cumulative Omicron cases detected in the State of Hawaii.

Variants Being Monitored in the State of Hawaii

Variant	Lineage	Area of emergence	Cumulative cases detected	Earliest specimen collection date	Most recent specimen collection date
Alpha			754		
	B.1.1.7	United Kingdom	703	21 Jan 2021	14 Aug 2021
	Q.3	United States	49	21 Mar 2021	02 Sep 2021
	Q.4	South Africa/Botswana	2	Apr 2021	Apr 2021
Beta	B.1.351	South Africa	19	16 Feb 2021	22 Sep 2021
Gamma			185		
	P.1.10	United States	81	24 Apr 2021	14 Jul 2021
	P.1	Brazil	52	24 Mar 2021	21 Jul 2021
	P.1.12	Peru	20	21 Mar 2021	28 Apr 2021
	P.1.17	United States/Mexico	18	29 Mar 2021	21 Jul 2021
	P.1.13	United States	14	03 May 2021	07 Jun 2021
Delta			4831		
	AY.103	United States	1028	21 May 2021	12 Jan 2022
	AY.44	United States	950	07 Jun 2021	19 Jan 2022
	Other AY.*	Various	709	01 Aug 2021	Sep 2021
	AY.3	United States	496	28 Jun 2021	02 Jan 2022
	AY.25	United States	432	21 Jun 2021	20 Jan 2022
	AY.54	United States	312	23 Jun 2021	21 Nov 2021
	AY.100	South Africa/Botswana	172	17 Jul 2021	12 Dec 2021
	AY.25.1	South Africa/Botswana	160	08 Jul 2021	27 Jan 2022
	AY.1	Europe	134	30 Jun 2021	30 Nov 2021
	AY.119	United States	134	06 Jul 2021	21 Dec 2021
	AY.26	United States/Mexico	117	07 Jun 2021	28 Dec 2021
	AY.117	United States	97	15 Jul 2021	17 Dec 2021
	B.1.617.2	India	90	28 May 2021	16 Dec 2021
Epsilon			774		
	B.1.429	California	715	31 Dec 2020	03 Jun 2021
	B.1.427	California	59	07 Dec 2020	05 Jun 2021
Zeta	P.2	Brazil	2	Feb 2021	Feb 2021
Iota	B.1.526	New York	128	06 Feb 2021	23 Jul 2021
Mu			53		
	B.1.621	Columbia	43	03 Jun 2021	17 Sep 2021
	B.1.621.1	United States	10	27 May 2021	11 Aug 2021

Table Notes:

- Lineage “Other AY.*” represents an aggregation of different AY.* lineages in which each alone accounts for <2% of the cumulative Delta cases detected in the State of Hawaii.

Honolulu County

Total variants identified




Figure Notes:

- The graph shows the total number of variants detected in Honolulu County in each 2-week interval ending on the date shown (date represents when the specimen was collected from a patient).
- Omicron (B.1.1.529 + BA.* + XE) is a *Variant of Concern*.
- Delta (B.1.617.2 + AY.*) is a *Variant Being Monitored*.
- The gray bar graph (top) shows the percentage of PCR-positive cases from each 2-week time interval that were sequenced.
- The color bar graph (bottom) shows the number of viruses sequenced (one virus equates to one PCR-positive case) from each 2-week time interval.
- SARS-CoV-2 genome sequencing may not be a random sample of all cases. This graph does not estimate prevalence in the population.
- Sequencing can be performed on stored patient specimens at any time, so these numbers may change as additional specimens are sequenced.

Estimated proportions of variants circulating in Honolulu County




Figure Notes:

- The graph shows biweekly percentage estimates of SARS-CoV-2 variants circulating in Honolulu County, grouped in two-week intervals (based on the date of specimen collection).
- Not all positive SARS-CoV-2 specimens are sequenced and sequenced specimens are not a random selection of all COVID-19 cases in Honolulu County. This graph has been generated only counting specimens that were selected randomly for the purpose of surveillance of community variants, to avoid over-representing the specimens that were selected for investigations of clusters.
- Sequencing of certain specimens can be delayed for technical reasons. Therefore, the dataset used for this report is dynamic and specimens that are newly sequenced will be added to the dataset as sequencing occurs. Because of this, trends based on historical data can change over time.
- Omicron (BA.1*) is an aggregation of BA.1 and its sub-lineages (i.e., BA.1.1, BA.1.15, etc.).
- Omicron (BA.2*) is an aggregation of BA.2, its sub-lineages (i.e., BA.2.3, BA.2.9, etc.), and XE; excludes BA.2.12.1.

Variants of Concern in Honolulu County

Variant	Lineage	Area of emergence	Cumulative cases detected	Earliest specimen collection date	Most recent specimen collection date
Omicron			2957		
	BA.1.1	South Africa/Botswana	684	27 Nov 2021	11 May 2022
	BA.2	South Africa/Botswana	575	27 Jan 2022	03 Jun 2022
	BA.2.3	Philippines	504	30 Jan 2022	02 Jun 2022
	BA.2.12.1	United States/Canada	431	21 Mar 2022	04 Jun 2022
	BA.1.1.2	Japan	300	06 Dec 2021	31 Mar 2022
	BA.1.15	United States	164	30 Nov 2021	07 Apr 2022
	BA.2.9	Europe	59	14 Mar 2022	01 Jun 2022
	Other BA.*	Various	49	02 Apr 2022	Mar 2022
	BA.1	South Africa/Botswana	43	07 Dec 2021	14 Mar 2022
	BA.1.1.18	United States	35	10 Dec 2021	30 Mar 2022
	BA.2.18	United Kingdom	23	05 Apr 2022	03 Jun 2022
	BA.2.13	Europe	14	25 Apr 2022	23 May 2022
	BA.1.20	United States	12	14 Dec 2021	05 Feb 2022
	BA.2.10.1	Singapore	11	03 Mar 2022	25 Apr 2022
	BA.2.38	India	10	21 Mar 2022	09 May 2022
	XE	United Kingdom	9	26 Mar 2022	31 May 2022
	BA.1.17.2	Europe	8	14 Dec 2021	23 Feb 2022
	BA.5	South Africa	8	14 May 2022	29 May 2022
	BA.2.32	Indonesia	7	08 Mar 2022	25 Apr 2022
	BA.2.7	United States	7	28 Mar 2022	21 May 2022
	BA.4	South Africa	4	20 May 2022	27 May 2022

Table Notes:

- Lineage “Other BA.*” represents an aggregation of different BA.* lineages, except BA.4, in which each alone accounts for <0.2% of the cumulative Omicron cases detected in Honolulu County.

Variants Being Monitored in Honolulu County

Variant	Lineage	Area of emergence	Cumulative cases detected	Earliest specimen collection date	Most recent specimen collection date
Alpha			477		
	B.1.1.7	United Kingdom	444	21 Jan 2021	14 Aug 2021
	Q.3	United States	33	03 Apr 2021	02 Sep 2021
Beta	B.1.351	South Africa	16	16 Feb 2021	22 Sep 2021
Gamma			94		
	P.1.10	United States	50	24 Apr 2021	13 Jul 2021
	P.1	Brazil	15	24 Mar 2021	11 Jul 2021
	P.1.13	United States	14	03 May 2021	07 Jun 2021
	P.1.17	United States/Mexico	14	29 Mar 2021	20 Jul 2021
	P.1.12	Peru	1	Apr 2021	Apr 2021
Delta			2182		
	AY.103	United States	582	20 Jun 2021	18 Dec 2021
	AY.44	United States	363	07 Jun 2021	22 Dec 2021
	Other AY.*	Various	244	01 Aug 2021	Sep 2021
	AY.54	United States	212	23 Jun 2021	21 Nov 2021
	AY.3	United States	202	28 Jun 2021	17 Dec 2021
	AY.25	United States	162	21 Jun 2021	04 Jan 2022
	AY.100	South Africa/Botswana	81	23 Jul 2021	12 Dec 2021
	AY.1	Europe	74	30 Jun 2021	30 Nov 2021
	AY.26	United States/Mexico	60	07 Jun 2021	21 Nov 2021
	AY.25.1	South Africa/Botswana	57	18 Jul 2021	11 Dec 2021
	AY.117	United States	50	15 Jul 2021	11 Nov 2021
	AY.119	United States	48	06 Jul 2021	13 Dec 2021
	B.1.617.2	India	47	28 May 2021	25 Oct 2021
Epsilon			346		
	B.1.429	California	315	05 Jan 2021	29 May 2021
	B.1.427	California	31	07 Jan 2021	05 Jun 2021
Zeta	P.2	Brazil	2	Feb 2021	Feb 2021
Iota	B.1.526	New York	26	08 Feb 2021	23 Jul 2021
Mu			21		
	B.1.621	Columbia	18	03 Jun 2021	28 Jul 2021
	B.1.621.1	United States	3	May 2021	Aug 2021

Table Notes:

- Lineage “Other AY.*” represents an aggregation of different AY.* lineages in which each alone accounts for <2% of the cumulative Delta cases detected in Honolulu County.

Maui County

Total variants identified




Figure Notes:

- The graph shows the total number of variants detected in Maui County in each 2-week interval ending on the date shown (date represents when the specimen was collected from a patient).
- Omicron (B.1.1.529 + BA.* + XE) is a *Variant of Concern*.
- The gray bar graph (top) shows the percentage of PCR-positive cases from each 2-week time interval that were sequenced.
- The color bar graph (bottom) shows the number of viruses sequenced (one virus equates to one PCR-positive case) from each 2-week time interval.
- SARS-CoV-2 genome sequencing may not be a random sample of all cases. This graph does not estimate prevalence in the population.
- Sequencing can be performed on stored patient specimens at any time, so these numbers may change as additional specimens are sequenced.

Estimated proportions of variants circulating in Maui County




Figure Notes:

- The graph shows biweekly percentage estimates of SARS-CoV-2 variants circulating in Maui County, grouped in two-week intervals (based on the date of specimen collection).
- Not all positive SARS-CoV-2 specimens are sequenced and sequenced specimens are not a random selection of all COVID-19 cases in Maui County. This graph has been generated only counting specimens that were selected randomly for the purpose of surveillance of community variants, to avoid over-representing the specimens that were selected for investigations of clusters.
- Sequencing of certain specimens can be delayed for technical reasons. Therefore, the dataset used for this report is dynamic and specimens that are newly sequenced will be added to the dataset as sequencing occurs. Because of this, trends based on historical data can change over time.
- Omicron (BA.1*) is an aggregation of BA.1 and its sub-lineages (i.e., BA.1.1, BA.1.15, etc.).
- Omicron (BA.2*) is an aggregation of BA.2, its sub-lineages (i.e., BA.2.3, BA.2.9, etc.), and XE; excludes BA.2.12.1.

Variants of Concern in Maui County

Variant	Lineage	Area of emergence	Cumulative cases detected	Earliest specimen collection date	Most recent specimen collection date
Omicron			954		
	BA.1.1	South Africa/Botswana	255	16 Dec 2021	25 May 2022
	BA.2	South Africa/Botswana	228	13 Feb 2022	30 May 2022
	BA.2.12.1	United States/Canada	159	09 Apr 2022	03 Jun 2022
	BA.1.15	United States	86	14 Dec 2021	09 Apr 2022
	BA.2.3	Philippines	68	28 Jan 2022	24 May 2022
	BA.1	South Africa/Botswana	26	17 Dec 2021	25 Feb 2022
	BA.1.1.2	Japan	23	13 Dec 2021	15 Feb 2022
	BA.1.1.18	United States	17	30 Dec 2021	22 Feb 2022
	BA.2.29	Japan	13	08 Apr 2022	12 May 2022
	BA.2.31	Israel	13	27 Apr 2022	24 May 2022
	BA.1.20	United States	9	19 Dec 2021	28 Feb 2022
	BA.2.18	United Kingdom	9	21 Mar 2022	20 May 2022
	BA.2.9	Europe	8	08 Mar 2022	18 May 2022
	BA.2.10	Asia	8	22 Feb 2022	14 Apr 2022
	BA.2.38	India	4	06 Apr 2022	18 Apr 2022
	BA.1.17.2	Europe	3	Dec 2021	Feb 2022
	BA.2.10.1	Singapore	3	Mar 2022	Apr 2022
	BA.2.37	United Kingdom	3	Mar 2022	Apr 2022
	BA.1.10	United Kingdom	2	Dec 2021	Dec 2021
	BA.2.7	United States	2	Apr 2022	May 2022
	BA.1.1.16	Canada/United States	2	Feb 2022	Feb 2022
	BA.4	South Africa	2	May 2022	May 2022
	BA.1.18	Europe/North America	2	Dec 2021	Dec 2021
	BA.1.15.2	United States	2	Jan 2022	Feb 2022
	BA.2.26	United States	1	Apr 2022	Apr 2022
	BA.2.22	United Kingdom	1	Apr 2022	Apr 2022
	BA.2.20	Canada	1	Apr 2022	Apr 2022
	BA.1.1.17	United States	1	Jan 2022	Jan 2022
	BA.2.6	France	1	May 2022	May 2022
	BA.2.9.2	United States	1	May 2022	May 2022
	BA.2.21	Canada/United States	1	Apr 2022	Apr 2022

Variants Being Monitored in Maui County

Variant	Lineage	Area of emergence	Cumulative cases detected	Earliest specimen collection date	Most recent specimen collection date
Alpha			41		
	B.1.1.7	United Kingdom	39	01 Mar 2021	11 Jul 2021
	Q.3	United States	2	Apr 2021	May 2021
Beta	B.1.351	South Africa	1	Jul 2021	Jul 2021
Gamma			42		
	P.1	Brazil	22	18 Apr 2021	20 Jun 2021
	P.1.12	Peru	19	21 Mar 2021	28 Apr 2021
	P.1.10	United States	1	May 2021	May 2021
Delta			370		
	AY.103	United States	80	13 Jul 2021	23 Dec 2021
	AY.44	United States	59	27 Jun 2021	27 Dec 2021
	AY.3	United States	47	19 Jul 2021	01 Dec 2021
	Other AY.*	Various	37	09 Jul 2021	Sep 2021
	AY.47	United States	31	19 Aug 2021	07 Dec 2021
	AY.100	South Africa/Botswana	19	22 Jul 2021	18 Nov 2021
	AY.13	United States	18	21 Jul 2021	13 Sep 2021
	AY.25	United States	16	19 Aug 2021	19 Dec 2021
	AY.2	United States	14	07 Jun 2021	06 Aug 2021
	AY.26	United States/Mexico	13	03 Aug 2021	28 Oct 2021
	AY.25.1	South Africa/Botswana	12	24 Jul 2021	10 Dec 2021
	AY.122	South Africa/Botswana	11	19 Jul 2021	01 Oct 2021
	AY.39	United States	11	25 Sep 2021	31 Oct 2021
	B.1.617.2	India	2	Dec 2021	Dec 2021
Epsilon			272		
	B.1.429	California	265	08 Jan 2021	18 May 2021
	B.1.427	California	7	27 Apr 2021	10 May 2021
Iota	B.1.526	New York	16	12 Mar 2021	25 Jun 2021

Table Notes:

- Lineage “Other AY.*” represents an aggregation of different AY.* lineages in which each alone accounts for <2% of the cumulative Delta cases detected in Maui County.

Hawaii County

Total variants identified




Figure Notes:

- The graph shows the total number of variants detected in Hawaii County in each 2-week interval ending on the date shown (date represents when the specimen was collected from a patient).
- Omicron (B.1.1.529 + BA.* + XE) is a *Variant of Concern*.
- The gray bar graph (top) shows the percentage of PCR-positive cases from each 2-week time interval that were sequenced.
- The color bar graph (bottom) shows the number of viruses sequenced (one virus equates to one PCR-positive case) from each 2-week time interval.
- SARS-CoV-2 genome sequencing may not be a random sample of all cases. This graph does not estimate prevalence in the population.
- Sequencing can be performed on stored patient specimens at any time, so these numbers may change as additional specimens are sequenced.

Estimated proportions of variants circulating in Hawaii County




Figure Notes:

- The graph shows biweekly percentage estimates of SARS-CoV-2 variants circulating in Hawaii County, grouped in two-week intervals (based on the date of specimen collection).
- Not all positive SARS-CoV-2 specimens are sequenced and sequenced specimens are not a random selection of all COVID-19 cases in Hawaii County. This graph has been generated only counting specimens that were selected randomly for the purpose of surveillance of community variants, to avoid over-representing the specimens that were selected for investigations of clusters.
- Sequencing of certain specimens can be delayed for technical reasons. Therefore, the dataset used for this report is dynamic and specimens that are newly sequenced will be added to the dataset as sequencing occurs. Because of this, trends based on historical data can change over time.
- Omicron (BA.1*) is an aggregation of BA.1 and its sub-lineages (i.e., BA.1.1, BA.1.15, etc.).
- Omicron (BA.2*) is an aggregation of BA.2, its sub-lineages (i.e., BA.2.3, BA.2.9, etc.), and XE; excludes BA.2.12.1.

Variants of Concern in Hawaii County

Variant	Lineage	Area of emergence	Cumulative cases detected	Earliest specimen collection date	Most recent specimen collection date
Omicron			713		
	BA.1.1	South Africa/Botswana	159	13 Dec 2021	09 May 2022
	BA.2	South Africa/Botswana	156	19 Jan 2022	04 Jun 2022
	BA.2.3	Philippines	93	30 Jan 2022	01 Jun 2022
	BA.2.12.1	United States/Canada	91	13 Apr 2022	03 Jun 2022
	BA.1.1.2	Japan	64	10 Dec 2021	01 Mar 2022
	BA.1	South Africa/Botswana	25	17 Dec 2021	02 Mar 2022
	BA.1.15	United States	24	10 Dec 2021	22 Mar 2022
	BA.2.9	Europe	20	28 Mar 2022	16 May 2022
	BA.2.18	United Kingdom	14	24 Apr 2022	26 May 2022
	BA.1.1.16	Canada/United States	8	28 Jan 2022	06 Mar 2022
	BA.1.1.18	United States	7	15 Dec 2021	02 Mar 2022
	BA.1.17.2	Europe	7	30 Dec 2021	25 Feb 2022
	BA.1.20	United States	5	21 Dec 2021	03 Jan 2022
	BA.2.10.1	Singapore	4	23 Feb 2022	13 Apr 2022
	BA.5	South Africa	4	03 May 2022	21 May 2022
	BA.2.3.4	Canada	4	19 Apr 2022	03 May 2022
	BA.2.32	Indonesia	4	06 May 2022	16 May 2022
	BA.2.7	United States	3	Apr 2022	May 2022
	BA.2.10	Asia	3	Apr 2022	May 2022
	BA.2.17	Vietnam/Japan	3	Mar 2022	Mar 2022
	BA.2.21	Canada/United States	3	May 2022	May 2022
	BA.2.12	North America/Europe	2	Feb 2022	Mar 2022
	BA.1.17	Europe	2	Jan 2022	Mar 2022
	BA.1.18	Europe/North America	1	Dec 2021	Dec 2021
	BA.2.8	United Kingdom	1	May 2022	May 2022
	BA.2.23	United Kingdom	1	May 2022	May 2022
	BA.2.36	Europe	1	May 2022	May 2022
	BA.2.31	Israel	1	May 2022	May 2022
	BA.1.1.14	Europe	1	Dec 2021	Dec 2021
	BA.2.13	Europe	1	May 2022	May 2022
	XE	United Kingdom	1	Apr 2022	Apr 2022

Variants Being Monitored in Hawaii County

Variant	Lineage	Area of emergence	Cumulative cases detected	Earliest specimen collection date	Most recent specimen collection date
Alpha			66		
	B.1.1.7	United Kingdom	55	22 Feb 2021	16 Jul 2021
	Q.3	United States	11	21 Mar 2021	20 Aug 2021
Gamma			16		
	P.1.10	United States	12	27 May 2021	13 Jun 2021
	P.1	Brazil	3	May 2021	Jun 2021
	P.1.17	United States/Mexico	1	Jun 2021	Jun 2021
Delta			737		
	AY.44	United States	209	11 Jun 2021	16 Dec 2021
	AY.3	United States	109	31 Jul 2021	16 Dec 2021
	AY.103	United States	91	21 May 2021	16 Dec 2021
	Other AY.*	Various	74	01 Jul 2021	Oct 2021
	AY.25	United States	52	19 Jul 2021	09 Nov 2021
	AY.100	South Africa/Botswana	41	17 Jul 2021	12 Nov 2021
	AY.46.4	United States	27	21 Jun 2021	06 Dec 2021
	AY.118	United States	26	08 Jul 2021	27 Sep 2021
	AY.25.1	South Africa/Botswana	26	11 Aug 2021	17 Dec 2021
	AY.117	United States	24	15 Aug 2021	17 Nov 2021
	AY.119	United States	24	03 Aug 2021	24 Nov 2021
	AY.26	United States/Mexico	16	24 Jun 2021	03 Dec 2021
	AY.122	South Africa/Botswana	15	19 Jul 2021	22 Sep 2021
	B.1.617.2	India	3	Jul 2021	Nov 2021
Epsilon			44		
	B.1.429	California	38	25 Jan 2021	26 May 2021
	B.1.427	California	6	04 Feb 2021	17 May 2021
Iota	B.1.526	New York	73	06 Feb 2021	07 Jul 2021
Mu	B.1.621	Columbia	1	Jun 2021	Jun 2021

Table Notes:

- Lineage “Other AY.*” represents an aggregation of different AY.* lineages in which each alone accounts for <2% of the cumulative Delta cases detected in Hawaii County.

Kauai County

Total variants identified




Figure Notes:

- The graph shows the total number of variants detected in Kauai County in each 2-week interval ending on the date shown (date represents when the specimen was collected from a patient).
- Omicron (B.1.1.529 + BA.* + XE) is a *Variant of Concern*.
- The gray bar graph (top) shows the percentage of PCR-positive cases from each 2-week time interval that were sequenced.
- The color bar graph (bottom) shows the number of viruses sequenced (one virus equates to one PCR-positive case) from each 2-week time interval.
- SARS-CoV-2 genome sequencing may not be a random sample of all cases. This graph does not estimate prevalence in the population.
- Sequencing can be performed on stored patient specimens at any time, so these numbers may change as additional specimens are sequenced.

Estimated proportions of variants circulating in Kauai County




Figure Notes:

- The graph shows biweekly percentage estimates of SARS-CoV-2 variants circulating in Kauai County, grouped in two-week intervals (based on the date of specimen collection).
- Not all positive SARS-CoV-2 specimens are sequenced and sequenced specimens are not a random selection of all COVID-19 cases in Kauai County. This graph has been generated only counting specimens that were selected randomly for the purpose of surveillance of community variants, to avoid over-representing the specimens that were selected for investigations of clusters.
- Sequencing of certain specimens can be delayed for technical reasons. Therefore, the dataset used for this report is dynamic and specimens that are newly sequenced will be added to the dataset as sequencing occurs. Because of this, trends based on historical data can change over time.
- Omicron (BA.1*) is an aggregation of BA.1 and its sub-lineages (i.e., BA.1.1, BA.1.15, etc.).
- Omicron (BA.2*) is an aggregation of BA.2, its sub-lineages (i.e., BA.2.3, BA.2.9, etc.), and XE; excludes BA.2.12.1.

Variants of Concern in Kauai County

Variant	Lineage	Area of emergence	Cumulative cases detected	Earliest specimen collection date	Most recent specimen collection date
Omicron			260		
	BA.2	South Africa/Botswana	83	17 Mar 2022	31 May 2022
	BA.1.1	South Africa/Botswana	63	15 Dec 2021	19 Apr 2022
	BA.2.12.1	United States/Canada	27	30 Mar 2022	02 Jun 2022
	BA.2.3	Philippines	21	09 Mar 2022	24 May 2022
	BA.1.15	United States	13	28 Dec 2021	23 Mar 2022
	BA.1	South Africa/Botswana	12	16 Dec 2021	07 Feb 2022
	BA.1.1.18	United States	8	28 Dec 2021	21 Mar 2022
	BA.1.1.2	Japan	8	15 Dec 2021	09 Feb 2022
	BA.2.9	Europe	5	17 Mar 2022	09 May 2022
	BA.2.18	United Kingdom	5	24 Apr 2022	19 May 2022
	BA.2.29	Japan	2	Apr 2022	Apr 2022
	BA.2.7	United States	2	May 2022	May 2022
	BA.1.18	Europe/North America	2	Dec 2021	Jan 2022
	BA.2.10	Asia	2	Apr 2022	May 2022
	BA.2.38	India	1	Apr 2022	Apr 2022
	BA.2.1	United Kingdom	1	May 2022	May 2022
	BA.2.13	Europe	1	May 2022	May 2022
	BA.1.20	United States	1	Jan 2022	Jan 2022
	BA.1.17.2	Europe	1	Feb 2022	Feb 2022
	BA.1.1.14	Europe	1	Feb 2022	Feb 2022
	BA.5	South Africa	1	Jun 2022	Jun 2022

Variants Being Monitored in Kauai County

Variant	Lineage	Area of emergence	Cumulative cases detected	Earliest specimen collection date	Most recent specimen collection date
Alpha			19		
	B.1.1.7	United Kingdom	18	05 Apr 2021	13 Jul 2021
	Q.4	South Africa/Botswana	1	Apr 2021	Apr 2021
Gamma			2		
	P.1	Brazil	1	May 2021	May 2021
	P.1.10	United States	1	May 2021	May 2021
Delta			233		
	AY.25	United States	70	16 Jul 2021	10 Nov 2021
	AY.119	United States	35	28 Aug 2021	21 Dec 2021
	AY.44	United States	34	01 Jul 2021	21 Dec 2021
	Other AY.*	Various	25	Aug 2021	Sep 2021
	AY.47	United States	19	27 Jul 2021	29 Nov 2021
	AY.103	United States	16	01 Aug 2021	15 Dec 2021
	AY.1	Europe	10	09 Aug 2021	23 Aug 2021
	AY.3	United States	8	30 Sep 2021	15 Dec 2021
	AY.54	United States	8	06 Jul 2021	09 Nov 2021
	AY.67	South Africa/Botswana	6	06 Jun 2021	29 Jun 2021
Epsilon	B.1.617.2	India	2	Sep 2021	Oct 2021
			30		
	B.1.429	California	28	07 Jan 2021	08 May 2021
	B.1.427	California	2	Apr 2021	Apr 2021
Iota	B.1.526	New York	1	Apr 2021	Apr 2021
Mu	B.1.621	Columbia	1	Jul 2021	Jul 2021

Table Notes:

- Lineage “Other AY.*” represents an aggregation of different AY.* lineages in which each alone accounts for <2% of the cumulative Delta cases detected in Kauai County.