State of Hawaii SARS-CoV-2 Sequencing and Variant Report Hawaii Department of Health

2022-10-25

Contents

Introduction	3
Acknowledgements	4
Summary and key notes	5
	6 6
XZ, XAC, XAP, XAS, and XBB lineages)	6 6
Beta variant (B.1.351 and B.1.351.* lineages)	6 7 7
Delta variant (B.1.617.2 and AY.* lineages)	7 7
	7 7 7
	8
Total variants identified	8 9 10
Total variants identified 1 Estimated proportions of variants circulating in Honolulu County 1 Variants of Concern in Honolulu County 1	.2 12 13 14
	.6
Estimated proportions of variants circulating in Maui County	16 17 18
Hawaii County 2	20
Estimated proportions of variants circulating in Hawaii County	20 21 22 23
	24
Estimated proportions of variants circulating in Kauai County	24 25 26 27

Introduction

Whole genome sequencing (WGS) involves a set of laboratory methods used to determine the full genome sequence of an organism or virus, which in the case of Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2), the virus that causes Coronavirus Disease 2019 (COVID-19), is approximately 30,000 letters, each letter a nucleotide code denoted as A, C, G, or T.

The genome sequence of a virus can reveal mutations that make it unique. Mutations are changes in a genome sequence (usually one-letter changes) that occur naturally over time.

Collecting the genome sequences of virus specimens can reveal information about the relatedness of viruses and the similarities shared among groups of viruses. Groups of same-species viruses that share a set of genome mutations are referred to as a lineage.

Scientists compare viral genomes to better understand virus transmission, how viruses can spread from person to person. Sequencing also allows public health officials to monitor viruses involved in outbreaks, characterize outbreaks, detect clusters of cases, and monitor new lineages. Novel mutations can emerge with new lineages and scientists refer to these new lineages as emerging variants.

Some of these variants are classified by the Centers for Disease Control and Prevention (CDC) as a Variant of Concern (VOC) and others as Variants Being Monitored (VBM), because of their attributes, which, for example, can be increased transmissibility, decreased neutralization by antibodies generated during previous infection or vaccination, and/or increased severity of disease. The CDC has extensive information about SARS-CoV-2 variant classification (https://www.cdc.gov/coronavirus/2019-ncov/cases-updates/variant-surveillance/variant-info.html), which is updated as new evidence becomes available.

Sequencing can only be performed on specimens that contain SARS-CoV-2 RNA, which means only specimens used for molecular tests (such as PCR) can be included. Therefore, this report is limited to confirmed PCR-positives only. The genomes that are sequenced and compared are those of the virus, not humans.

Sequencing can be performed on stored specimens at any time. Therefore, the dataset used for this report is dynamic and batches of stored specimens that are newly sequenced will be added to the dataset as sequencing occurs. Because of this, trends based on historical data can change over time.

In February 2021, State Laboratories Division, Hawaii Department of Health increased sequencing efforts done on PCR-positive specimens to improve the State's ability to detect new variants and conduct genomic surveillance of SARS-CoV-2. According to the CDC, approximately 6% of PCR-positive cases in the State of Hawaii have been sequenced since testing began (https://covid.cdc.gov/covid-data-tracker/#published-sars-cov-2-sequences).

Acknowledgements

Institution	Program/partner	Count	Percent
State Laboratories Division		14984	73.025%
Centers for Disease Control and Prevention	National SARS-CoV-2 Strain Surveillance (NS3)	412	2.008%
	Quest Diagnostics Incorporated	1246	6.072%
	Laboratory Corporation of America	1521	7.413%
	Aegis Sciences Corporation	379	1.847%
	Helix/Illumina	174	0.848%
	Infinity Biologix	16	0.078%
	Mako Medical	20	0.097%
	Fulgent Genetics	8	0.039%
Tripler Army Medical Center		682	3.324%
University of Hawaii		861	4.196%
Aegis Sciences Corporation [*]		216	1.053%
Total		20519	100.000%

Total 14984

Total 861

Total

216

148

This report integrates genomes sequenced since Jan 1, 2021 by:

County distribution of genomes sequenced by:

State Laboratories I	Division (since J	an 1, 2021)		
Honolulu County	Maui County	Hawaii County	Kauai County	unknown
8435	3062	2399	739	349
University of Hawai	i (Jan 22, 2022 -	– Aug 23, 2022)		
				-
Honolulu County	Maui County	Hawaii County	Kauai County	unknown
Honolulu County 508	Maui County 67	Hawaii County 221	Kauai County 17	unknown 48
	67	221	<i>.</i>	

3

11

Notes:

54

• In support of genomic surveillance efforts, PCR-positive specimens are routinely provided to State Laboratories Division (SLD) by commercial laboratories that operate throughout the State of Hawaii, including Clinical Labs of Hawaii (CLH), Diagnostic Laboratory Services, Inc. (DLS), Kaiser Permanente Hawaii, and S&G Labs Hawaii. The specimens are then processed, sequenced, and analyzed by the Laboratory Preparedness and Response Branch (LPRB) within SLD.

0

- In coordination with SLD, the School of Life Sciences, University of Hawaii at Manoa provided sequencing • and analysis services for 861 PCR-positive specimens processed by the LPRB.
- As of September 14, 2022, Aegis Sciences Corporation* reports SARS-CoV-2 genome sequences independently of the Centers for Disease Control and Prevention (CDC).
- County information is not provided for specimens sequenced by CDC programs/partners and by Tripler • Army Medical Center (TAMC).

Summary and key notes

- This report reflects sequenced PCR-positive specimens only, and not all PCR-positive cases in the State of Hawaii are sequenced.
- Each successfully sequenced specimen produces one consensus SARS-CoV-2 genome sequence that is further analyzed to determine the variant.
- State Laboratories Division reported 384 additional SARS-CoV-2 genome sequences since the previous report was generated (10/12/2022).
- CDC programs/partners reported 78 additional SARS-CoV-2 genome sequences from the State of Hawaii since the previous report was generated (10/12/2022).
- TAMC reported one additional SARS-CoV-2 genome sequence from the State of Hawaii since the previous report was generated (10/12/2022).
- SARS-CoV-2 variant nomenclature is defined by a World Health Organization (WHO) label (letters of the Greek Alphabet, e.g., Alpha, Beta, Gamma, Delta, etc.), Phylogenetic Assignment of Named Global Outbreak (PANGO) lineage (alphabetical prefix and a numerical suffix), and/or Nextstrain clade (year of emergence followed by the next available letter in the alphabet, e.g., 20A, 20B, etc.). In this report, variant counts are reported using the WHO label and PANGO lineage nomenclatures only.
- For PANGO lineage nomenclature, 'the numerical suffix has three hierarchical levels (primary, secondary, and tertiary). Each full stop (period or dot) within the numerical suffix represents "descendant of". Descendants of lineages with tertiary suffixes are assigned to the next available alphabetical prefix, in alphabetical order.' More details can be found at https://www.pango.network/the-pango-nomenclature-system/statement-of-nomenclature-rules/.
- In this report, an asterisk (*) following a lineage designation (e.g., BA.5*) signifies inclusion of the respective lineage (e.g., BA.5) and its sub-lineages (or descendants, e.g., BA.5.*) unless otherwise specified. An asterisk following the dot of a lineage designation (e.g., BA.*, BA.5.*, BA.5.1.*) signifies only inclusion of lineages with an additional numerical suffix that follows the respective dot.
- Lineages called using pangolin v4.1.3, pangolin-data v1.15.1, and usher v0.5.4.
- Omicron has represented 100% of the variants circulating in the State of Hawaii since 1/28/2022.
- Classifications of Omicron lineages are in flux. The parent lineage of Omicron is B.1.1.529; primary sub-lineages of B.1.1.529 have BA.* designations (e.g., BA.1, BA.2, BA.5). Primary sub-lineages of BA.2.12.1, BA.2.75.1, BA.2.75.3, BA.2.75.6, BA.5.1.10, BA.5.2.1, and BA.5.3.1 have BG.*, BL.*, BM.*, BY.*, BK.*, BF.*, and BE.* designations, respectively. The primary sub-lineage of BE.1.1.1 has the BQ.* designation. XE, XZ, XAC, XAP, XAS, and XBB are Omicron recombinants; a recombinant forms when the genomes of two lineages (infecting a person simultaneously) combine during the viral replication process, producing a variant that is distinct from both parent lineages.
- BA.5* (BA.5 and its sub-lineages) continues to decline proportionally but remains dominant, representing ~82% of the sequenced specimens collected in the State of Hawaii from 9/25/2022 to 10/8/2022.
- BQ.1.1, which has recently been detected locally, is currently the fastest growing lineage nationally with more than a 2-fold estimated growth advantage over BA.5*, and contains at least 3 additional immune escape mutations within the spike protein compared to BA.5*.
- Estimated proportions of variants circulating may not be truly representative in counties with low sequencing numbers (e.g., Kauai County).

Variants of Concern (VOC) and Variants Being Monitored (VBM)

Evidence to date shows that vaccination generally leads to milder cases, not requiring hospitalization, for all Variants of Concern and Variants Being Monitored that are described here, even if the efficacy of antibodies is diminished against some of these variants compared to the original version of the virus.

Also, none of these variants are classified as a "Variant of High Consequence (VOHC)", according to CDC variant categories (https://www.cdc.gov/coronavirus/2019-ncov/variants/variant-info.html#Consequence), which is a category that would imply a variant has the ability to evade diagnosis, significantly reduce the vaccines' effectiveness and protection against severe disease, significantly reduce susceptibility to treatments, or lead to more severe disease and increased hospitalizations.

Variants of Concern detected in the State of Hawaii

Omicron variant (B.1.1.529, BA.*, BE.*, BF.*, BG.*, BK.*, BL.*, BM.*, BQ.*, BY.*, XE, XZ, XAC, XAP, XAS, and XBB lineages)

B.1.1.529 was reported to the WHO on November 24, 2021 and first detected in specimens collected on November 11, 2021 in Botswana and on November 14, 2021 in South Africa. The WHO labeled B.1.1.529 "Omicron" and classified it as a VOC on November 26, 2021. The United States designated Omicron as a VOC on November 30, 2021 and reported its first case on December 1, 2021. Omicron contains more changes in the spike protein than have been observed in other variants. Several of these mutations, including S477N, N501Y, and E484K, have been associated with increased infectivity and decreased neutralizing activity of monoclonal antibodies and convalescent sera. Evidence indicates that Omicron spreads more easily than previous variants. More information can be found at https://www.cdc.gov/coronavirus/2019-ncov/variants/omicron-variant.html.

B.1.1.529 is the parent lineage of Omicron; primary sub-lineages of the Omicron parent lineage have BA.* designations (e.g., BA.1, BA.2, BA.5). Primary sub-lineages of BA.2.12.1, BA.2.75.1, BA.2.75.3, BA.2.75.6, BA.5.1.10, BA.5.2.1, and BA.5.3.1 have BG.*, BL.*, BM.*, BY.*, BK.*, BF.*, and BE.* designations, respectively. The primary sub-lineage of BE.1.1.1 has the BQ.* designation. SARS-CoV-2 recombinants that receive a PANGO classification have an X* lineage designation. XE, XZ, XAC, and XAP are Omicron recombinants with portions of their genomes derived from BA.1* and BA.2*. XAS is an Omicron recombinant with portions of its genome derived from BA.2* and BA.5*, and XBB is an Omicron recombinant with portions of its genome derived from BJ.1 (a primary sub-lineage of BA.2.10.1) and BM.1.1.1. Omicron lineage families correspond to Nextstrain clades 21M, 21K, 21L, 22A, 22B, 22C, 22D, and 22E (https://ncov-clades-schema.vercel.app/).

Due to acquired mutations within the spike protein, each successively dominant Omicron lineage (BA.1^{*}, BA.2^{*}, BA.2.12.1^{*}, and BA.5^{*}) has been increasingly better at evading antibodies elicited by vaccinations and/or prior infections. Notable information regarding each of their defining mutations can be found at https://covariants.org/variants (21K, 21L, 22C, and 22B, respectively).

Variants Being Monitored detected in the State of Hawaii

Alpha variant (B.1.1.7 and Q.* lineages)

B.1.1.7 was first identified in the United Kingdom and the WHO labeled it "Alpha" on May 31, 2021. This variant contains the N501Y mutation and a short deletion in the spike protein. Upon its emergence, the Alpha variant was shown to be significantly more transmissible (\sim 50%) than the original SARS-CoV-2 virus. The Alpha variant corresponds to Nextstrain clade 20I.

Beta variant (B.1.351 and B.1.351.* lineages)

B.1.351 was first identified in South Africa and the WHO labeled it "Beta" on May 31, 2021. This variant has the E484K spike mutation, which is associated with increased evasion of antibodies elicited by vaccination or previous infection by earlier variants. The Beta variant corresponds to Nextstrain clade 20H.

Gamma variant (P.1 and P.1.* lineages)

P.1 was first identified in Brazil and the WHO labeled it "Gamma" on May 31, 2021. This variant also contains the N501Y mutation, like B.1.1.7, but not the deletion in the spike protein. The Gamma variant corresponds to Nextstrain clade 20J.

Delta variant (B.1.617.2 and AY.* lineages)

B.1.617.2 was first identified in India and the WHO labeled it "Delta" on May 31, 2021. This variant contains the L452R mutation in the spike protein, which has been shown to escape neutralization by monoclonal antibodies and some convalescent sera, as well as a few additional spike mutations predicted to have functional consequences (e.g. T478K).

B.1.617.2 is the parent lineage of Delta; sub-lineages of Delta have the AY.* designation. Delta lineage families correspond to Nextstrain clades 21A, 21I, and 21J.

Epsilon variant (B.1.429 and B.1.427)

The closely related lineages, B.1.429 and B.1.427, were first identified in California and designated initially as CA VUI1. The WHO labeled them "Epsilon" on May 31, 2021. This variant is associated with increased viral shedding compared to the original SARS-CoV-2 virus, and contains the L452R mutation in the spike protein, which has been shown to escape neutralization by monoclonal antibodies and some convalescent sera. The Epsilon variant corresponds to Nextstrain clade 21C.

Zeta variant (P.2)

P.2 was first identified in Brazil and contains a spike mutation (E484K), which is also present in B.1.351, that can potentially make it less responsive to antibodies. The Zeta variant corresponds to Nextstrain clade 20B/S.484K.


Iota variant (B.1.526)

B.1.526 was first identified in New York and was classified by the CDC as a VBM due to indications of increased transmissibility. The WHO labeled it "Iota" on May 31, 2021. Some of the genomes (but not all) of this variant contain the E484K mutation. The Iota variant corresponds to Nextstrain clade 21F.

Mu variant (B.1.621 and B.1.621.1)


Lineage B.1.621 was first identified in Columbia in January 2021 and has a couple of mutations in common with the Beta (B.1.351) and Gamma (P.1) variants, which have been associated with increased transmissibility (N501Y) and a level of decreased vaccine efficiency (E484K). The MU variant corresponds to Nextstrain clade 21H.

State of Hawaii



Total variants identified

- The graph shows the total number of variants by lineage identified in the State of Hawaii in each 2-week interval ending on the date shown (date represents when the specimen was collected from a patient).
- All lineages identified within the overall time period shown belong to the Omicron variant.
- Upper (gray) bar graph shows the percentage of PCR-positive cases from each 2-week time interval that were sequenced.
- Lower (color) bar graph shows the number of viruses sequenced (one sequenced virus equates to one PCR-positive case) from each 2-week time interval.
- SARS-CoV-2 genome sequencing may not be a random sample of all cases. This graph does not estimate prevalence in the population.
- Sequencing can be performed on stored patient specimens at any time, so these numbers may change as additional specimens are sequenced.



Estimated proportions of variants circulating in the State of Hawaii

- The graph shows biweekly percentage estimates of SARS-CoV-2 variants circulating in the State of Hawaii, grouped in 2-week intervals (based on the date of specimen collection); percentage estimates are rounded to the nearest whole number.
- Not all positive SARS-CoV-2 specimens are sequenced and sequenced specimens are not a random selection of all COVID-19 cases in the State of Hawaii. This graph has been generated only counting specimens that were selected randomly for the purpose of surveillance.
- The last 2-week interval numbers will most likely change, as a number of specimens that are currently being processed will be added.
- Sequencing of certain specimens can be delayed for technical reasons. Therefore, the dataset used for this report is dynamic and specimens that are newly sequenced will be added to the dataset as sequencing occurs. Because of this, trends based on historical data can change over time.
- Omicron BA.1* includes BA.1 and its sub-lineages (BA.1.*).
- Omicron BA.2* includes BA.2, its sub-lineages (BA.2.*), XE, XZ, XAC, and XAP; excludes BA.2.3.20, BA.2.12.1*, and BA.2.75*.
- Omicron BA.2.12.1* includes BA.2.12.1 and its sub-lineages (BG.*).
- Omicron BA.2.75* includes BA.2.75 and its sub-lineages (BA.2.75.*, BL.*, BM.*, BY.*).
- Omicron BA.4* includes BA.4 and its sub-lineages (BA.4.*); excludes BA.4.6.
- Omicron BA.5* includes BA.5, its sub-lineages (BA.5.*, BE.*, BF.*, BK.*), and XAS; excludes BF.7.
- Omicron BQ.1* includes BQ.1 and its sub-lineages (BQ.1.*); excludes BQ.1.1.
- Omicron XBB* includes XBB and its sub-lineages (XBB.*).

Variant	Lineage	Area of emergence	Cumulative sequences	Earliest specimen collection date	Most recent specimen collection date
Omicron			12567		
	BA.1.1	South Africa/Botswana	1740	27 Nov 2021	19 Jun 2022
	BA.2.12.1	USA/Canada	1614	21 Mar 2022	29 Sep 2022
	BA.2	South Africa/Botswana	1573	18 Jan 2022	06 Sep 2022
	Other	Various	1484	$01 \ \text{Feb} \ 2022$	31 May 2022
	BA.5.2.1	South Africa/UK/USA	901	31 May 2022	09 Oct 2022
	BA.2.3	Philippines	839	06 Jan 2022	22 Jul 2022
	BA.1.1.2	Japan	569	06 Dec 2021	01 Apr 2022
	BG.5	USA	533	$09 { m Apr} 2022$	21 Aug 2022
	BA.1.15	USA	426	30 Nov 2021	09 Apr 2022
	BA.5.2	South Africa/UK/USA	386	17 Jun 2022	11 Oct 2022
	BA.5.5	USA	357	03 May 2022	08 Oct 2022
	BA.5.1	Portugal	344	07 Jun 2022	11 Oct 2022
	BA.5.6	USA	232	21 May 2022	10 Oct 2022
	BA.2.3.17	USA	208	25 Jan 2022	15 Jun 2022
	BA.1	South Africa/Botswana	187	07 Dec 2021	11 Apr 2022
	BA.4.1	South Africa	168	04 May 2022	30 Sep 2022
	BA.2.9	Europe	164	25 Jan 2022	25 Jul 2022
	BA.5.1.1	USA	160	14 May 2022	06 Oct 2022
	BA.4.6	USA/UK/Denmark	130	21 Jun 2022	11 Oct 2022
	BF.5	Israel	125	16 Jun 2022	07 Oct 2022
	BA.1.1.18	USA	95	10 Dec 2021	30 Mar 2022
	BA.2.18	UK	93	21 Mar 2022	11 Jul 2022
	BF.10	USA	74	05 Jun 2022	07 Oct 2022
	BA.5.1.23	Spain	73	23 Jun 2022	04 Oct 2022
	BA.5.2.9	USA	72	20 Jun 2022	08 Oct 2022
	BQ.1	Nigeria	7	16 Sep 2022	04 Oct 2022
	BQ.1.1	Global	4	27 Sep 2022	07 Oct 2022
	XBB.1	Bangladesh/Singapore	4	30 Sep 2022	07 Oct 2022
	BQ.1.2	Global	2	03 Oct 2022	03 Oct 2022
	BQ.1.5	Global	2	02 Oct 2022	09 Oct 2022
	BQ.1.3	Global	1	15 Sep 2022	15 Sep 2022

Variants of Concern in the State of Hawaii

Table Notes:

• Lineage "Other" represents an aggregation of different Omicron lineages in which each alone accounts for <0.4% of the cumulative Omicron sequences; excludes BQ.1* and XBB* lineages.

Variant	Lineage	Area of emergence	Cumulative sequences	Earliest specimen collection date	Most recent specimen collection date
Alpha			797		
r	B.1.1.7	UK	743	21 Jan 2021	14 Aug 2021
	Q.3	USA	52	21 Mar 2021	02 Sep 2021
	Q.4	South Africa/Botswana	2	28 Apr 2021	29 Apr 2021
Beta	B.1.351	South Africa	20	16 Feb 2021	22 Sep 2021
Gamma			191		
	P.1.10	USA	84	24 Apr 2021	14 Jul 2021
	P.1	Brazil	54	24 Mar 2021	21 Jul 2021
	P.1.12	Peru	20	21 Mar 2021	28 Apr 2021
	P.1.17	USA/Mexico	18	29 Mar 2021	21 Jul 2021
Delta	P.1.13	USA	$\frac{15}{5622}$	03 May 2021	28 Jun 2021
Dena	AY.103	USA	1228	21 May 2021	12 Jan 2022
	AY.44	USA	1228	07 Jun 2021	12 Jan 2022 19 Jan 2022
	Other	Various	612	01 Aug 2021	30 Jul 2021
	AY.*	Various	012	01 Aug 2021	50 Jul 2021
	AY.3	USA	562	24 Jun 2021	02 Jan 2022
	AY.25	USA	532	21 Jun 2021	20 Jan 2022
	AY.54	USA	367	23 Jun 2021	21 Nov 2021
	AY.100	South Africa/Botswana	197	17 Jul 2021	12 Dec 2021
	AY.25.1	South Africa/Botswana	196	08 Jul 2021	27 Jan 2022
	AY.1	Europe	186	30 Jun 2021	30 Nov 2021
	AY.119	USA	136	06 Jul 2021	21 Dec 2021
	AY.26	USA/Mexico	132	07 Jun 2021	28 Dec 2021
	AY.117	USA	128	15 Jul 2021	17 Dec 2021
	AY.122	South Africa/Botswana	107	09 Jul 2021	26 Nov 2021
	B.1.617.2	India	102	28 May 2021	16 Dec 2021
	AY.47	USA	98	21 Jul 2021	07 Dec 2021
Epsilon			781		
-	B.1.429	USA	722	31 Dec 2020	03 Jun 2021
	B.1.427	USA	59	07 Dec 2020	05 Jun 2021
Zeta	P.2	Brazil	2	06 Feb 2021	08 Feb 2021
Iota	B.1.526	USA	128	06 Feb 2021	23 Jul 2021
Mu			58		
	B.1.621	Columbia	45	03 Jun 2021	17 Sep 2021
	B.1.621.1	USA	13	27 May 2021	24 Aug 2021


Variants Being Monitored in the State of Hawaii

Table Notes:


• Lineage "Other AY.*" represents an aggregation of different AY.* lineages in which each alone accounts for <2% of the cumulative Delta sequences from the State of Hawaii.

Honolulu County

Total variants identified



- The graph shows the total number of variants by lineage identified in Honolulu County in each 2-week interval ending on the date shown (date represents when the specimen was collected from a patient).
- All lineages identified within the overall time period shown belong to the Omicron variant.
- Upper (gray) bar graph shows the percentage of PCR-positive cases from each 2-week time interval that were sequenced.
- Lower (color) bar graph shows the number of viruses sequenced (one sequenced virus equates to one PCR-positive case) from each 2-week time interval.
- SARS-CoV-2 genome sequencing may not be a random sample of all cases. This graph does not estimate prevalence in the population.
- Sequencing can be performed on stored patient specimens at any time, so these numbers may change as additional specimens are sequenced.



Estimated proportions of variants circulating in Honolulu County

- The graph shows biweekly percentage estimates of SARS-CoV-2 variants circulating in Honolulu County, grouped in 2-week intervals (based on the date of specimen collection); percentage estimates are rounded to the nearest whole number.
- Not all positive SARS-CoV-2 specimens are sequenced and sequenced specimens are not a random selection of all COVID-19 cases in Honolulu County. This graph has been generated only counting specimens that were selected randomly for the purpose of surveillance of community variants.
- Sequencing of certain specimens can be delayed for technical reasons. Therefore, the dataset used for this report is dynamic and specimens that are newly sequenced will be added to the dataset as sequencing occurs. Because of this, trends based on historical data can change over time.
- Omicron BA.2* includes BA.2, its sub-lineages (BA.2.*), XE, and XAP; excludes BA.2.3.20, BA.2.12.1*, and BA.2.75*.
- Omicron BA.2.12.1* includes BA.2.12.1 and its sub-lineages (BG.*).
- Omicron BA.2.75* includes BA.2.75 and its sub-lineages (BA.2.75.*, BL.*, BM.*).
- Omicron BA.4* includes BA.4 and its sub-lineages (BA.4.*); excludes BA.4.6.
- Omicron BA.5* includes BA.5 and its sub-lineages (BA.5.*, BE.*, BF.*, BK.*); excludes BF.7.
- Omicron BQ.1* includes BQ.1 and its sub-lineages (BQ.1.*); excludes BQ.1.1.
- Omicron XBB* includes XBB and its sub-lineages (XBB.*).

Variant	Lineage	Area of emergence	Cumulative sequences	Earliest specimen collection date	Most recent specimen collection date
Omicron			5224		
	BA.1.1	South Africa/Botswana	683	27 Nov 2021	11 May 2022
	BA.2	South Africa/Botswana	623	27 Jan 2022	20 Jul 2022
	Other	Various	531	01 Aug 2022	31 May 2022
	BA.2.12.1	USA/Canada	504	$21~{\rm Mar}~2022$	29 Sep 2022
	BA.5.2.1	South Africa/UK/USA	387	05 Jun 2022	08 Oct 2022
	BA.2.3	Philippines	367	30 Jan 2022	24 Jun 2022
	BG.5	USA	303	09 Apr 2022	21 Aug 2022
	BA.1.1.2	Japan	300	06 Dec 2021	31 Mar 2022
	BA.5.2	South Africa/UK/USA	192	22 Jun 2022	11 Oct 2022
	BA.5.5	USA	165	29 May 2022	05 Oct 2022
	BA.1.15	USA	164	30 Nov 2021	07 Apr 2022
	BA.5.1	Portugal	141	11 Jun 2022	11 Oct 2022
	BA.2.3.17	USA	121	05 Feb 2022	02 Jun 2022
	BA.5.6	USA	121	18 Jun 2022	01 Oct 2022
	BA.4.1	South Africa	81	22 May 2022	04 Sep 2022
	BA.5.1.1	USA	74	14 May 2022	06 Oct 2022
	BA.2.9	Europe	70	14 Mar 2022	29 Jun 2022
	BA.4.6	USA/UK/Denmark	61	26 Jun 2022	11 Oct 2022
	BF.5	Israel	46	17 Jun 2022	06 Oct 2022
	BA.1	South Africa/Botswana	43	07 Dec 2021	14 Mar 2022
	BA.5.1.23	Spain	36	24 Jun 2022	16 Aug 2022
	BA.1.1.18	USA	35	10 Dec 2021	30 Mar 2022
	BA.2.3.14	South Korea	30	04 Apr 2022	30 May 2022
	BA.2.18	UK	29	$12~{\rm Apr}~2022$	01 Jul 2022
	BE.1.1	Germany	29	18 Jun 2022	03 Oct 2022
	BF.10	USA	27	09 Jul 2022	07 Oct 2022
	BA.5.2.21	Indonesia	25	11 Jul 2022	06 Oct 2022
	BF.7	UK/Belgium/Denmarl	25	19 Jul 2022	12 Oct 2022
	BQ.1	Nigeria	3	$23~{\rm Sep}~2022$	04 Oct 2022
	XBB.1	Bangladesh/Singapore	3	04 Oct 2022	07 Oct 2022
	BQ.1.2	Global	2	03 Oct 2022	03 Oct 2022
	BQ.1.1	Global	1	07 Oct 2022	07 Oct 2022
	BQ.1.3	Global	1	$15 { m Sep} \ 2022$	$15 { m Sep} \ 2022$
	BQ.1.5	Global	1	02 Oct 2022	02 Oct 2022

Variants of Concern in Honolulu County

Table Notes:

• Lineage "Other" represents an aggregation of different Omicron lineages in which each alone accounts for <0.4% of the cumulative Omicron sequences; excludes BQ.1* and XBB* lineages.

Variant	Lineage	Area of emergence	Cumulative sequences	Earliest specimen collection date	Most recent specimen collection date
Alpha			514		
1	B.1.1.7	UK	481	21 Jan 2021	14 Aug 2021
	Q.3	USA	33	03 Apr 2021	02 Sep 2021
Beta	B.1.351	South Africa	17	16 Feb 2021	22 Sep 2021
Gamma			98		-
	P.1.10	USA	53	24 Apr 2021	13 Jul 2021
	P.1	Brazil	15	24 Mar 2021	11 Jul 2021
	P.1.13	USA	15	03 May 2021	28 Jun 2021
	P.1.17	USA/Mexico	14	29 Mar 2021	20 Jul 2021
	P.1.12	Peru	1	03 Apr 2021	03 Apr 2021
Delta			2635		
	AY.103	USA	732	20 Jun 2021	18 Dec 2021
	AY.44	USA	417	07 Jun 2021	22 Dec 2021
	Other AY.*	Various	283	01 Aug 2021	30 Nov 2021
	AY.54	USA	259	23 Jun 2021	21 Nov 2021
	AY.3	USA	237	24 Jun 2021	17 Dec 2021
	AY.25	USA	200	21 Jun 2021	04 Jan 2022
	AY.1	Europe	107	30 Jun 2021	30 Nov 2021
	AY.100	South Africa/Botswana	88	23 Jul 2021	12 Dec 2021
	AY.117	USA	74	15 Jul 2021	11 Nov 2021
	AY.26	USA/Mexico	69	07 Jun 2021	21 Nov 2021
	AY.25.1	South Africa/Botswana	64	18 Jul 2021	11 Dec 2021
	B.1.617.2	India	55	28 May 2021	25 Oct 2021
	AY.119	USA	50	06 Jul 2021	13 Dec 2021
Epsilon			351		
	B.1.429	USA	320	05 Jan 2021	29 May 2021
	B.1.427	USA	31	07 Jan 2021	05 Jun 2021
Zeta	P.2	Brazil	2	06 Feb 2021	08 Feb 2021
Iota	B.1.526	USA	26	08 Feb 2021	23 Jul 2021
Mu			25		
	B.1.621	Columbia	19	03 Jun 2021	28 Jul 2021
	B.1.621.1	USA	6	27 May 2021	24 Aug 2021

Variants Being Monitored in Honolulu County

Table Notes:


• Lineage "Other AY.*" represents an aggregation of different AY.* lineages in which each alone accounts for <2% of the cumulative Delta sequences from Honolulu County.

Maui County

50-49% 40% 40 Lineage % of PCR-positive cases sequenced 38% BA.1.1 BA.4.1 BA.5.3 34% BA.2 BA.4.1.6 BA.5.3.1 30-BA.2.3 BA.4.2 BA.5.5 BA.2.3.17 BA.4.4 BA.5.5.2 BA.2.3.20 BA.4.5 BA.5.6 BA.2.6 BA.4.6 BA.5.6.2 20-20% 19% BA.2.7 BA.5 BA.5.8 19% BA.2.9 BA.5.1 BA.5.10 16% 15% BA.2.13 BA.5.1.1 BA.5.10.1 12% BA.2.18 BA.5.1.2 BE.1 10-BA.2.31 BA.5.1.3 BE.1.1 BE.1.1.1 BA.2.31.1 BA.5.1.6 BA.2.37 BA.5.1.7 BE.2 0-BA.2.38 BA.5.1.19 BE.3 BA.5.1.22 BE.4 BA.2.41 200-BA.2.42 BA.5.1.23 BF.1 BA.2.47 BA.5.1.25 BF.5 BA.2.48 BA.5.2 BF.7 BA.2.52 BA.5.2.1 BF.8 150-BA.2.65 BA.5.2.6 BF.9 Number of viruses sequencec BA.2.73 BA.5.2.8 BF.10 XE BA.5.2.9 BF.11 BA.2.12.1 BA.5.2.19 BF.21 BG.2 BA.5.2.20 BF.26 100 BG.4 BA.5.2.21 BF.27 BG.5 BA.5.2.22 BF.28 BA.2.75.1 BA.5.2.27 BK.1 BA.2.75.2 BA.5.2.28 BQ.1 BY.1 BA.5.2.31 50 BA.4 BA.5.2.33 0-2022-07-16 2022-07-30 2022-08-13 2022-09-10 2022-09-24 2022-06-18 2022-07-02 2022-08-21 2022-10-08 2022-06-04 2-week period ending on date

Total variants identified

- The graph shows the total number of variants by lineage identified in Maui County in each 2-week interval ending on the date shown (date represents when the specimen was collected from a patient).
- All lineages identified within the overall time period shown belong to the Omicron variant.
- Upper (gray) bar graph shows the percentage of PCR-positive cases from each 2-week time interval that were sequenced.
- Lower (color) bar graph shows the number of viruses sequenced (one sequenced virus equates to one PCR-positive case) from each 2-week time interval.
- SARS-CoV-2 genome sequencing may not be a random sample of all cases. This graph does not estimate prevalence in the population.
- Sequencing can be performed on stored patient specimens at any time, so these numbers may change as additional specimens are sequenced.



Estimated proportions of variants circulating in Maui County

- The graph shows biweekly percentage estimates of SARS-CoV-2 variants circulating in Maui County, grouped in 2-week intervals (based on the date of specimen collection); percentage estimates are rounded to the nearest whole number.
- Not all positive SARS-CoV-2 specimens are sequenced and sequenced specimens are not a random selection of all COVID-19 cases in Maui County. This graph has been generated only counting specimens that were selected randomly for the purpose of surveillance of community variants.
- Sequencing of certain specimens can be delayed for technical reasons. Therefore, the dataset used for this report is dynamic and specimens that are newly sequenced will be added to the dataset as sequencing occurs. Because of this, trends based on historical data can change over time.
- Omicron BA.1* includes BA.1 and its sub-lineages (BA.1.*).
- Omicron BA.2* includes BA.2, its sub-lineages (BA.2.*), XE, and XAC; excludes BA.2.3.20, BA.2.12.1*, and BA.2.75*.
- Omicron BA.2.12.1* includes BA.2.12.1 and its sub-lineages (BG.*).
- Omicron BA.2.75* includes BA.2.75 and its sub-lineages (BA.2.75.*, BY.*)
- Omicron BA.4* includes BA.4 and its sub-lineages (BA.4.*); excludes BA.4.6.
- Omicron BA.5* includes BA.5 and its sub-lineages (BA.5.*, BE.*, BF.*, BK.*); excludes BF.7.
- Omicron BQ.1* includes BQ.1 and its sub-lineages (BQ.1.*); excludes BQ.1.1.

Variant	Lineage	Area of emergence	Cumulative sequences	Earliest specimen collection date	Most recent specimen collection date
Omicron			2326		
	BA.2.12.1	USA/Canada	413	06 Apr 2022	26 Aug 2022
	BA.1.1	South Africa/Botswana	257	16 Dec 2021	19 Jun 2022
	BA.2	South Africa/Botswana	254	13 Feb 2022	12 Jul 2022
	BA.5.2.1	South Africa/UK/USA	238	31 May 2022	07 Oct 2022
	Other	Various	227	01 Aug 2022	31 May 2022
	BA.5.2	South Africa/UK/USA	95	19 Jun 2022	07 Oct 2022
	BA.5.5	USA	94	02 Jun 2022	$07 { m Sep } 2022$
	BA.5.1	Portugal	91	17 Jun 2022	03 Oct 2022
	BA.1.15	USA	86	14 Dec 2021	09 Apr 2022
	BA.2.3	Philippines	72	28 Jan 2022	14 Jul 2022
	BF.5	Israel	48	16 Jul 2022	05 Oct 2022
	BA.5.6	USA	41	27 May 2022	24 Sep 2022
	BA.5.2.9	USA	39	01 Jul 2022	06 Sep 2022
	BG.5	USA	37	19 Apr 2022	16 Jul 2022
	BA.4.1	South Africa	35	09 May 2022	06 Sep 2022
	BA.4.6	USA/UK/Denmark	30	29 Jun 2022	30 Sep 2022
	BA.5.1.23	Spain	28	23 Jun 2022	04 Oct 2022
	BA.1	South Africa/Botswana	26	17 Dec 2021	25 Feb 2022
	BA.5.1.1	USA	26	17 Jun 2022	17 Aug 2022
	BA.1.1.2	Japan	23	13 Dec 2021	15 Feb 2022
	BA.2.9	Europe	20	08 Mar 2022	25 Jul 2022
	BA.2.42	Australia	19	24 Apr 2022	23 Jun 2022
	BF.10	USA	18	23 Jun 2022	08 Sep 2022
	BA.1.1.18	USA	17	30 Dec 2021	22 Feb 2022
	BE.3	USA	17	28 Jun 2022	29 Sep 2022
	BA.2.29	Japan	15	08 Apr 2022	16 May 2022
	BA.2.18	UK	14	21 Mar 2022	11 Jul 2022
	BF.27	Singapore	12	05 Jul 2022	01 Aug 2022
	BA.5.8	USA/UK/Denmark	11	30 Jun 2022	23 Jul 2022
	BA.2.3.17	USA	10	21 Mar 2022	15 Jun 2022
	BA.2.31	Israel	10	02 May 2022	06 Jun 2022
	BQ.1	Nigeria	3	16 Sep 2022	03 Oct 2022

Variants of Concern in Maui County

Table Notes:

• Lineage "Other" represents an aggregation of different Omicron lineages in which each alone accounts for <0.4% of the cumulative Omicron sequences; excludes BQ.1* and XBB* lineages.

Variant	Lineage	Area of emergence	Cumulative sequences	Earliest specimen collection date	Most recent specimen collection date
Alpha			41		
1	B.1.1.7	UK	39	01 Mar 2021	11 Jul 2021
	Q.3	USA	2	03 Apr 2021	28 May 2021
Beta	B.1.351	South Africa	1	08 Jul 2021	08 Jul 2021
Gamma			43		
	P.1	Brazil	23	18 Apr 2021	20 Jun 2021
	P.1.12	Peru	19	21 Mar 2021	28 Apr 2021
	P.1.10	USA	1	10 May 2021	10 May 2021
Delta			434		
	AY.103	USA	95	13 Jul 2021	23 Dec 2021
	AY.44	USA	65	27 Jun 2021	27 Dec 2021
	AY.3	USA	60	19 Jul 2021	01 Dec 2021
	Other AY.*	Various	43	02 Aug 2021	30 Sep 2021
	AY.47	USA	31	19 Aug 2021	07 Dec 2021
	AY.100	South Africa/Botswana	23	22 Jul 2021	18 Nov 2021
	AY.13	USA	22	21 Jul 2021	13 Sep 2021
	AY.25	USA	18	12 Aug 2021	19 Dec 2021
	AY.25.1	South Africa/Botswana	18	24 Jul 2021	10 Dec 2021
	AY.122	South Africa/Botswana	16	19 Jul 2021	01 Oct 2021
	AY.26	USA/Mexico	16	03 Aug 2021	28 Oct 2021
	AY.2	USA	14	07 Jun 2021	06 Aug 2021
	AY.39	USA	11	25 Sep 2021	31 Oct 2021
	B.1.617.2	India	2	02 Dec 2021	16 Dec 2021
Epsilon			273		
	B.1.429	USA	266	08 Jan 2021	18 May 2021
	B.1.427	USA	7	27 Apr 2021	10 May 2021
Iota	B.1.526	USA	16	12 Mar 2021	25 Jun 2021


Variants Being Monitored in Maui County

Table Notes:


• Lineage "Other AY.*" represents an aggregation of different AY.* lineages in which each alone accounts for <2% of the cumulative Delta sequences from Maui County.

Hawaii County

Total variants identified



- The graph shows the total number of variants by lineage identified in Hawaii County in each 2-week interval ending on the date shown (date represents when the specimen was collected from a patient).
- All lineages identified within the overall time period shown belong to the Omicron variant.
- Upper (gray) bar graph shows the percentage of PCR-positive cases from each 2-week time interval that were sequenced.
- Lower (color) bar graph shows the number of viruses sequenced (one sequenced virus equates to one PCR-positive case) from each 2-week time interval.
- SARS-CoV-2 genome sequencing may not be a random sample of all cases. This graph does not estimate prevalence in the population.
- Sequencing can be performed on stored patient specimens at any time, so these numbers may change as additional specimens are sequenced.



Estimated proportions of variants circulating in Hawaii County

- The graph shows biweekly percentage estimates of SARS-CoV-2 variants circulating in Hawaii County, grouped in 2-week intervals (based on the date of specimen collection); percentage estimates are rounded to the nearest whole number.
- Not all positive SARS-CoV-2 specimens are sequenced and sequenced specimens are not a random selection of all COVID-19 cases in Hawaii County. This graph has been generated only counting specimens that were selected randomly for the purpose of surveillance of community variants.
- Sequencing of certain specimens can be delayed for technical reasons. Therefore, the dataset used for this report is dynamic and specimens that are newly sequenced will be added to the dataset as sequencing occurs. Because of this, trends based on historical data can change over time.
- Omicron BA.2* includes BA.2, its sub-lineages (BA.2.*), XE, XZ, and XAC; excludes BA.2.3.20, BA.2.12.1*, and BA.2.75*.
- Omicron BA.2.12.1* includes BA.2.12.1 and its sub-lineages (BG.*).
- Omicron BA.4* includes BA.4 and its sub-lineages (BA.4.*); excludes BA.4.6.
- Omicron BA.5* includes BA.5, its sub-lineages (BA.5.*, BE.*, BF.*, BK.*), and XAS; excludes BF.7.

Variant	Lineage	Area of emergence	Cumulative sequences	Earliest specimen collection date	Most recent specimen collection date
Omicron			1429		
	BA.2.12.1	USA/Canada	213	13 Apr 2022	12 Aug 2022
	BA.2	South Africa/Botswana	182	19 Jan 2022	06 Sep 2022
	BA.1.1	South Africa/Botswana	159	13 Dec 2021	09 May 2022
	BA.5.2.1	South Africa/UK/USA	142	12 Jun 2022	07 Oct 2022
	Other	Various	140	01 Aug 2022	31 May 2022
	BA.2.3	Philippines	90	30 Jan 2022	22 Jul 2022
	BA.1.1.2	Japan	64	10 Dec 2021	01 Mar 2022
	BA.5.1	Portugal	45	07 Jun 2022	10 Oct 2022
	BA.5.1.1	USA	41	14 May 2022	28 Sep 2022
	BA.5.2	South Africa/UK/USA	38	17 Jun 2022	07 Oct 2022
	BA.5.6	USA	32	21 May 2022	28 Sep 2022
	BA.5.5	USA	27	03 May 2022	12 Sep 2022
	BG.5	USA	26	13 Apr 2022	12 Jul 2022
	BA.1	South Africa/Botswana	25	17 Dec 2021	02 Mar 2022
	BA.1.15	USA	24	10 Dec 2021	22 Mar 2022
	BA.2.9	Europe	21	28 Mar 2022	03 Jul 2022
	BA.2.18	UK	19	24 Apr 2022	24 Jun 2022
	BA.4.1	South Africa	16	16 Jun 2022	01 Sep 2022
	BF.5	Israel	15	07 Jul 2022	01 Oct 2022
	BA.4.6	USA/UK/Denmark	14	11 Jul 2022	26 Sep 2022
	BA.2.3.17	USA	12	24 Feb 2022	05 May 2022
	BE.1	South Africa/UK/Austria	11	22 Jul 2022	06 Oct 2022
	BF.10	USA	11	06 Jul 2022	06 Oct 2022
	BE.3	USA	10	08 Jun 2022	07 Aug 2022
	BA.1.1.16	Canada/USA	9	28 Jan 2022	27 Apr 2022
	BA.1.1.18	USA	7	15 Dec 2021	02 Mar 2022
	BA.1.17.2	Europe	7	30 Dec 2021	25 Feb 2022
	BA.2.32	Indonesia	7	06 May 2022	16 Jul 2022
	BA.5.2.21	Indonesia	7	25 Jul 2022	07 Oct 2022
	BA.5.2.9	USA	7	14 Aug 2022	08 Oct 2022
	BA.5.1.23	Spain	6	27 Jun 2022	03 Aug 2022
	BQ.1.1	Global	2	30 Sep 2022	07 Oct 2022

Variants of Concern in Hawaii County

Table Notes:

• Lineage "Other" represents an aggregation of different Omicron lineages in which each alone accounts for <0.4% of the cumulative Omicron sequences; excludes BQ.1* and XBB* lineages.

Variant	Lineage	Area of emergence	Cumulative sequences	Earliest specimen collection date	Most recent specimen collection date
Alpha			71		
1	B.1.1.7	UK	57	22 Feb 2021	22 Jul 2021
	Q.3	USA	14	21 Mar 2021	20 Aug 2021
Gamma			17		
	P.1.10	USA	12	27 May 2021	13 Jun 2021
	P.1	Brazil	4	10 May 2021	02 Jun 2021
	P.1.17	USA/Mexico	1	14 Jun 2021	14 Jun 2021
Delta			951		
	AY.44	USA	229	11 Jun 2021	16 Dec 2021
	AY.3	USA	123	31 Jul 2021	16 Dec 2021
	AY.103	USA	113	21 May 2021	16 Dec 2021
	AY.25	USA	108	19 Jul 2021	09 Nov 2021
	Other AY.*	Various	84	01 Jul 2021	30 Jul 2021
	AY.100	South Africa/Botswana	54	17 Jul 2021	12 Nov 2021
	AY.25.1	South Africa/Botswana	45	28 Jul 2021	17 Dec 2021
	AY.118	USA	41	08 Jul 2021	27 Sep 2021
	AY.117	USA	30	11 Aug 2021	17 Nov 2021
	AY.122	South Africa/Botswana	29	19 Jul 2021	22 Sep 2021
	AY.46.4	USA	27	21 Jun 2021	06 Dec 2021
	AY.119	USA	24	03 Aug 2021	24 Nov 2021
	AY.26	USA/Mexico	19	24 Jun 2021	03 Dec 2021
	AY.1	Europe	18	20 Jul 2021	03 Sep 2021
	B.1.617.2	India	7	06 Jul 2021	08 Nov 2021
Epsilon			44		
	B.1.429	USA	38	25 Jan 2021	26 May 2021
	B.1.427	USA	6	04 Feb 2021	17 May 2021
Iota	B.1.526	USA	73	06 Feb 2021	07 Jul 2021
Mu	B.1.621	Columbia	1	13 Jun 2021	13 Jun 2021


Variants Being Monitored in Hawaii County

Table Notes:


• Lineage "Other AY.*" represents an aggregation of different AY.* lineages in which each alone accounts for <2% of the cumulative Delta sequences from Hawaii County.

Kauai County

Total variants identified



- The graph shows the total number of variants by lineage identified in Kauai County in each 2-week interval ending on the date shown (date represents when the specimen was collected from a patient).
- All lineages identified within the overall time period shown belong to the Omicron variant.
- Upper (gray) bar graph shows the percentage of PCR-positive cases from each 2-week time interval that were sequenced.
- Lower (color) bar graph shows the number of viruses sequenced (one sequenced virus equates to one PCR-positive case) from each 2-week time interval.
- SARS-CoV-2 genome sequencing may not be a random sample of all cases. This graph does not estimate prevalence in the population.
- Sequencing can be performed on stored patient specimens at any time, so these numbers may change as additional specimens are sequenced.



Estimated proportions of variants circulating in Kauai County

- The graph shows biweekly percentage estimates of SARS-CoV-2 variants circulating in Kauai County, grouped in 2-week intervals (based on the date of specimen collection); percentage estimates are rounded to the nearest whole number.
- Not all positive SARS-CoV-2 specimens are sequenced and sequenced specimens are not a random selection of all COVID-19 cases in Kauai County. This graph has been generated only counting specimens that were selected randomly for the purpose of surveillance of community variants.
- Sequencing of certain specimens can be delayed for technical reasons. Therefore, the dataset used for this report is dynamic and specimens that are newly sequenced will be added to the dataset as sequencing occurs. Because of this, trends based on historical data can change over time.
- Omicron BA.2* includes BA.2, its sub-lineages (BA.2.*), and XAC; excludes BA.2.3.20, BA.2.12.1*, and BA.2.75*.
- Omicron BA.2.12.1* includes BA.2.12.1 and its sub-lineages (BG.*).
- Omicron BA.4* includes BA.4 and its sub-lineages (BA.4.*); excludes BA.4.6.
- Omicron BA.5* includes BA.5 and its sub-lineages (BA.5.*, BE.*, BF.*, BK.*); excludes BF.7.
- Omicron BQ.1* includes BQ.1 and its sub-lineages (BQ.1.*); excludes BQ.1.1.

Variant	Lineage	Area of emergence	Cumulative sequences	Earliest specimen collection date	Most recent specimen collection date
Omicron			453		
	BA.2	South Africa/Botswana	83	17 Mar 2022	26 Jul 2022
	BA.1.1	South Africa/Botswana	63	15 Dec 2021	19 Apr 2022
	BA.2.12.1	USA/Canada	62	30 Mar 2022	27 Jul 2022
	Other	Various	42	02 Jun 2022	26 Jan 2022
	BA.5.2.1	South Africa/UK/USA	27	03 Jun 2022	07 Oct 2022
	BA.2.3	Philippines	19	09 Mar 2022	24 May 2022
	BA.5.5	USA	18	13 Jun 2022	31 Aug 2022
	BA.5.1	Portugal	15	23 Jun 2022	03 Oct 2022
	BA.1.15	USA	13	28 Dec 2021	23 Mar 2022
	BA.1	South Africa/Botswana	12	16 Dec 2021	07 Feb 2022
	BA.5.2	South Africa/UK/USA	9	02 Jul 2022	07 Oct 2022
	XAC	USA/Canada	9	28 Apr 2022	14 Jun 2022
	BA.1.1.18	USA	8	28 Dec 2021	21 Mar 2022
	BA.1.1.2	Japan	8	15 Dec 2021	09 Feb 2022
	BA.2.18	UK	8	24 Apr 2022	16 Jun 2022
	BA.2.42	Australia	8	07 May 2022	02 Jul 2022
	BG.5	USA	7	03 May 2022	13 Jun 2022
	BA.4.1	South Africa	6	08 Jun 2022	26 Jun 2022
	BA.4.6	$\rm USA/\rm UK/\rm Denmark$	6	23 Aug 2022	30 Sep 2022
	BF.10	USA	6	13 Jun 2022	19 Aug 2022
	BA.2.9	Europe	5	17 Mar 2022	09 May 2022
	BA.5.2.9	USA	4	27 Jun 2022	30 Aug 2022
	BA.5.6	USA	4	16 Jun 2022	29 Aug 2022
	BF.5	Israel	4	21 Jun 2022	06 Oct 2022
	BA.2.1	UK	3	25 May 2022	20 Jun 2022
	BE.3	USA	3	15 Jun 2022	15 Sep 2022
	BQ.1	Nigeria	1	30 Sep 2022	30 Sep 2022

Variants of Concern in Kauai County

Table Notes:

• Lineage "Other" represents an aggregation of different Omicron lineages in which each alone accounts for <0.5% of the cumulative Omicron sequences; excludes BQ.1* and XBB* lineages.

Variant	Lineage	Area of emergence	Cumulative sequences	Earliest specimen collection date	Most recent specimen collection date
Alpha			20		
-	B.1.1.7	UK	19	05 Apr 2021	13 Jul 2021
	Q.4	South Africa/Botswana	1	28 Apr 2021	28 Apr 2021
Gamma			2		
	P.1	Brazil	1	31 May 2021	31 May 2021
	P.1.10	USA	1	14 May 2021	14 May 2021
Delta			246	v	v
	AY.25	USA	70	16 Jul 2021	10 Nov 2021
	AY.44	USA	37	29 Jun 2021	21 Dec 2021
	AY.119	USA	35	28 Aug 2021	21 Dec 2021
	Other AY.*	Various	25	03 Nov 2021	29 Nov 2021
	AY.47	USA	19	27 Jul 2021	29 Nov 2021
	AY.1	Europe	18	09 Aug 2021	30 Aug 2021
	AY.103	USA	17	01 Aug 2021	15 Dec 2021
	AY.3	USA	8	$30 { m Sep } 2021$	15 Dec 2021
	AY.54	USA	8	06 Jul 2021	09 Nov 2021
	AY.67	South Africa/Botswana	7	06 Jun 2021	30 Jun 2021
	B.1.617.2	India	2	21 Sep 2021	18 Oct 2021
Epsilon			31	-	
	B.1.429	USA	29	07 Jan 2021	08 May 2021
	B.1.427	USA	2	27 Apr 2021	27 Apr 2021
Iota	B.1.526	USA	1	14 Apr 2021	14 Apr 2021
Mu	B.1.621	Columbia	2	29 Jun 2021	06 Jul 2021

Variants Being Monitored in Kauai County

Table Notes:

• Lineage "Other AY.*" represents an aggregation of different AY.* lineages in which each alone accounts for <2% of the cumulative Delta sequences from Kauai County.