

Name:

Birthdate:

Age:

Today's Date:

## CHILD LEAD RISK QUESTIONS

Did the child have an elevated blood lead level in the past?	Yes	No
--	-----	----

Does the child have a sibling or playmate with lead poisoning?	Yes	No
--	-----	----

Does the child live in or regularly visit a place built before 1978?	Yes	No
--	-----	----

Is it being or was it recently remodeled or renovated?	Yes	No
--	-----	----

Does anyone who spends time with the child have a job or hobby in:		
- Automotive repair / car batteries	Yes	No
- Making ceramics or pottery using lead glaze	Yes	No
- Fishing sinkers / fishing activities / boat repair	Yes	No
- Painting / electrical / plumbing / soldering / welding	Yes	No
- Remodeling / renovation / building demolition	Yes	No

Does the child often eat, chew, or mouth dirt, paint chips, or other things that are not food?	Yes	No
--	-----	----

Does the family cook or serve food using ceramic dishes or pottery that may have a lead glaze (such as Chinese or Mexican pottery)?	Yes	No
---	-----	----

Does the family have a catchment water system?	Yes	No
--	-----	----



Lead risk assessment is recommended at ages 9-12 months and 2 years (or at 3-6 years if not previously done) and as risk level changes.

Medicaid/EPSTD requirement: Blood lead levels are required at ages 9-12 months and 2 years. A blood lead level should be done at 3-6 years of age if a level has never been done or risk level changes.

For children with other insurance coverage: Blood lead levels are recommended if there are risk factors. Health plans vary in their coverage of blood lead levels.