



Hawaii State Department of Health

Disease Investigation Branch

Oahu 586-4586, Maui 984-8213, Hilo 933-0912,
Kona 322-4877, Kauai 241-3563

Swimmer's Itch

What is swimmer's itch?

Swimmer's itch is a rash caused by an allergic reaction to certain microscopic parasites that infect birds and animals. These parasites are released from infected snails into fresh and salt water. The infection is found throughout the world and is more frequent during summer months.

How do you get it?

You get it by swimming or wading in waters contaminated with the parasite. The parasite larvae burrow into human skin and causes allergic reaction and rash. The risk of being infected is greater in shallow water by the shoreline with warm temperature. Although anyone can get swimmer's itch, children are infected more often since they tend to swim, wade, and play in the shallow water more than adults. Also, they are less likely to towel dry themselves when leaving the water. Person-to-person spread does not occur, since these larvae cannot develop in a human and die soon.

What are the symptoms of swimmer's itch?

First you may experience tingling, burning, or itching of the skin. Small reddish pimples appear within 12 hours. Pimples may develop into small blisters. Itching may last up to a week or more, but will go away gradually. Scratching the areas may result in secondary bacterial infection. Because swimmer's itch is caused by allergic reaction, the more often you are exposed to contaminated water, the more intense and immediate symptoms of swimmer's itch will be.

When do symptoms start?

The first symptoms can begin within a few minutes to days of exposure to the contaminated water.

What is the treatment for swimmer's itch?

Fact Sheets
Revised March 14, 2019
Disease Investigation Branch

Most cases of swimmer's itch do not require medical attention. If you have a rash, you may try the following for relief:

- Use corticosteroid cream
- Apply cool compresses to the affected areas
- Bathe in Epsom salts or baking soda
- Soak in colloidal oatmeal baths
- Apply baking soda paste to the rash (made by stirring water into baking soda until it reaches a paste-like consistency)
- Use an anti-itch lotion

If itching is severe, your doctor may recommend a prescription medication.

Although it is very difficult, try not to scratch. Scratching may cause the rash to become infected.

How do you keep from getting it?

- Do not swim in areas where swimmer's itch is a known problem or where signs have been posted warning of unsafe water.
- Do not swim near or wade in marshy areas where snails are commonly found.
- Towel dry or shower immediately after leaving the water.
- Use protective gears such as rubber wading boots as needed.
- Do not attract birds to areas where people are swimming.

For more information, see CDC's website.

https://www.cdc.gov/parasites/swimmersitch/health_professionals/index.html