

COVID-19 Vaccine Timing 2025-26 – Routine Schedule

Age	Vaccine	If unvaccinated:	If had any prior doses, give 2025-26 doses:
6 – 23 months	Moderna–Spikevax (pediatric)		<p>If 1 prior Moderna dose: 1</p> <p>If 1 prior Pfizer dose: 1 2</p> <p>If ≥2 prior Moderna/Pfizer doses: 1</p>
2 – 4 years ^{^^}	Moderna–Spikevax (pediatric)	1 Dose	<p>If ≥1 prior Moderna/Pfizer dose: 1</p>
5 – 11 years ^{^^}	Moderna–Spikevax (pediatric)	1 Dose	<p>If 1 or more prior doses (of any of the brands), then[^]: 1 Spikevax/Comirnaty</p>
	Pfizer–Comirnaty (pediatric)	1 Dose	
12+ years ^{^^}	Pfizer–Comirnaty (adol/adult)	1 Dose	<p>If 1 or more prior doses (of any of the brands), then[^]:</p> <p>Ages 12–64 years^{^^}: 1 Spikevax/Mnexspike/Comirnaty/Nuvaxovid</p> <p>Ages 65+ years: 1 2</p>
	Moderna–Spikevax (adol/adult)	1 Dose	
	Moderna–Mnexspike	1 Dose	
	Novavax–Nuvaxovid	1 Dose	

* An [8-week interval](#) may be preferable for some people, especially for males 12-39 years.

[^] Janssen (J & J) vaccine has been deauthorized. Follow schedule for 12+ years for any prior doses.

[§] Minimum interval 2 months, except for Mnexspike, the minimum interval is 3 months.

^{^^} Children 2-18 years old are recommended vaccination if they are in [certain risk groups](#), have never been vaccinated, or their parent/guardian desires their protection from COVID-19.

View [additional pediatric vaccination guidance](#), and [adult vaccination guidance](#).
 For storage and administration information, see [COVID-19 Product Guide](#).



COVID-19 Vaccine Timing 2025-26 if Moderately/Severely Immunocompromised

Age	Vaccine	If unvaccinated:	If had any prior doses, give 2025-26 doses:
6 months–4 years	Moderna–Spikevax (pediatric)		If prior doses of Moderna or Pfizer at 6m–4y: <div> 1 prior dose: 4 w 1 ≥4 w 2 </div> <div> 2 prior doses: ≥4 w 1 </div> <div> ≥3 prior doses**: ≥8 w 1 (for ages 5+ yrs, Pfizer dose is also OK) </div> <div>6 m^s</div> <div>Additional Dose(s)*</div>
	Moderna–Spikevax (pediatric)		
5–11 years	Pfizer–Comirnaty (pediatric)		<div> 1 prior dose: 3 w 1 ≥4 w 2 </div> <div> 2 prior doses: ≥4 w 1 </div> <div> ≥3 prior doses**: ≥8 w 1 </div> <div>6 m^s</div> <div>Additional Dose(s)*</div>
	Pfizer–Comirnaty (pediatric)		
12+ years	Pfizer–Comirnaty (adol/adult)		<div> 1 prior dose: 3 w 1 ≥4 w 2 </div> <div> 2 prior doses: ≥4 w 1 </div> <div> ≥3 prior doses**: ≥8 w 1 </div> <div>6 m^s</div> <div>Additional Dose(s)*</div>
	Moderna–Spikevax (adol/adult)		<div> 1 prior dose: 4 w 1 ≥4 w 2 </div> <div> 2 prior doses: ≥4 w 1 </div> <div> ≥3 prior doses**: ≥8 w 1 </div> <div>6 m^s</div> <div>Additional Dose(s)*</div>
	Moderna–Mnexspike		<div> 1 prior dose: 4 w 1 ≥4 w 2 </div> <div> 2 prior doses: ≥4 w 1 </div> <div> ≥3 prior doses**: ≥8 w 1 </div> <div>6 m^s</div> <div>Additional Dose(s)*</div>
	Novavax–Nuvaxovid		<div> 1 prior dose: ≥3 w 1 </div> <div> ≥2 prior doses**: ≥8 w 1 </div> <div>6 m^s</div> <div>Additional Dose(s)*</div>

* **Further doses** may be given under shared clinical decision-making at a minimum interval of 2 months.

** Ages 5–11 years may be given Moderna or Pfizer after ≥3 prior doses. Ages 12+ years may be given Spikevax, Comirnaty, Nuvaxovid, or Mnexspike.

§ Minimum interval 2 months, except for Mnexspike, the minimum interval is 3 months.

¶ If >8 weeks passed since the first Novavax dose, any 2025–26 COVID-19 vaccine (Spikevax/Comirnaty/Nuvaxovid/Mnexspike) may be given.

View [additional pediatric vaccination guidance](#), and [adult vaccination guidance](#).

For storage and administration information, see [COVID-19 Product Guide](#).

