

Antibiogram for N. gonorrhoeae Isolates in Hawaii from 2010-2019

| YEAR | Number of Isolates Tested | Percent Susceptible | | | | |
|------|---------------------------|---------------------|---------------|----------|-------------|--------------|
| | | Ceftriaxone | Ciprofloxacin | Cefixime | Cefpodoxime | Azithromycin |
| 2010 | 218 | 100 | 62.0 | 100 | 97.2 | 98.1 |
| 2011 | 189 | 100 | 70.5 | 100 | 96.8 | 99.5 |
| 2012 | 189 | 100 | 82.0 | 100 | 99.5 | 98.4 |
| 2013 | 157 | 100 | 86.2 | 98.7 | 100 | 99.4 |
| 2014 | 210 | 100 | --- | 95.7 | --- | 99.0 |
| 2015 | 221 | 100 | --- | 98.1 | --- | 95.0 |
| 2016 | 267 | 97.8 | --- | 98.1 | --- | 87.6 |
| 2017 | 260 | 100 | --- | 100 | --- | 82.7 |
| 2018 | 267 | 99.6 | --- | 99.6 | --- | 92.5 |
| 2019 | 215 | 99.5 | --- | 98.6 | --- | 96.7 |

Antimicrobial resistance criteria

Ceftriaxone Alert value $\geq 0.125 \mu\text{g/ml}$

Cefixime Alert value $\geq 0.250 \mu\text{g/ml}$

Azithromycin Alert value $\geq 2.0 \mu\text{g/ml}$

Data was created and compiled at the Hawaii Department of Health State Laboratories

This antibiogram supercedes previous versions. Previous data may have been adjusted to meet current interpretive guidelines.